

*19th Biennial Meeting of the ISACB
Vienna, Austria
October 5-8, 2024*



POSTER ABSTRACTS

ISACB 2024 ABSTRACT

Loss of NOTCH1 Promotes Impaired Vascular Smooth Muscle Cell Response to Hemodynamic Stress and Ascending Aortic Aneurysms

Authors: Ruth Ackah MD^{1,2}, Anupama Rao PhD², Michael Boutros BS³, Jun Yasuhara MD², Yukie Ueyama B.V.Sc², Ming-Tao Zhao PhD², Vidu Garg MD^{2,4}

1 Department of Surgery, The Ohio State University Wexner Medical Center, Columbus, OH, USA

2 Center for Cardiovascular Research and Heart Center, Nationwide Children's Hospital, Columbus Ohio, USA

3 Department of Pediatrics, The Ohio State University, Columbus, Ohio, USA

4 Department of Molecular Genetics, The Ohio State University, Columbus, Ohio, USA

Background/Objective: NOTCH1 is a critical cardiovascular developmental gene. Mutations in NOTCH1 are linked to BAV, Ascending Aortic Aneurysms (AscAA), and other of cardiovascular defects in humans. This study sought to investigate the potential mechanisms by which NOTCH1 haploinsufficiency produces the AscAA phenotype and the impact of NOTCH1 haploinsufficiency on vascular smooth muscle cell (VSMC) response to hemodynamic stress.

Methods: Three human induced pluripotent stem cell (iPSC) lines (control, NOTCH1^{+/-}, and NOTCH1^{-/-}) were generated using CRISPR/Cas9 genome editing and differentiated into VSMCs via second heart field (SHF) lineage. Differentiation of VSMC was performed by adding TGFβ and PDGF-BB to the culture media of SHF progenitor cells for ~12 days. Immunohistochemistry was performed to assess differentiation to SHF progenitors and differentiated VSMCs from each line were characterized for contractile SMC markers by immunofluorescence. Cells were exposed to cyclic stress over 24hrs and RNA collected for Bulk RNA-seq analysis.

Results: Immunohistochemistry staining revealed that expression of SHF cardiac progenitor population markers (ISL1) was not significantly different among control, NOTCH1^{+/-}, and NOTCH1^{-/-} mutants cell lines. However, expression of mature SMC-specific contractile proteins (SMA and MYH11) was significantly lower in both NOTCH1^{+/-}, and NOTCH1^{-/-} cells compared to control SMCs. Bulk RNA-seq analysis demonstrated a significant impairment of NOTCH1^{+/-} VSMCs to generate a canonical contractile VSMC response to mechanical stress compared to control. RNA expression profiles of NOTCH1^{+/-} VSMCs exposed to hemodynamic stress revealed a more pro-inflammatory/pro-apoptotic response and evidence of VSMC contractile to synthetic phenotype switching.

Conclusions: Our results indicate that loss of NOTCH1 in the human iPSC model did not affect differentiation of iPSCs to SHF progenitor cells but is crucial for the differentiation of contractile VSMCs from SHF progenitor cells. Mutations in NOTCH1 may predispose to the development of AscAA by affecting contractile VSMC differentiation from SHF progenitors and provokes an impaired VSMC response to stress.

Funding: This research was supported by the National Institutes of Health under the Ruth L. Kirschstein National Research Service Award (F32), grant number 1F32HL160059 from the National Heart, Lung, and Blood Institute (NHLBI).

Financial disclosures: None

Alarmin S100A8/A9 drives histone methylation-based epigenetic alterations in blood- and bone marrow-derived leukocytes in acute myocardial infarction

Alexandra-Gela Lazar¹, Mihaela-Loredana Vlad¹, Razvan Gheorghita Mares², Simona-Adriana Manea¹, Mihai Bogdan Preda¹, Maya Simionescu¹, Adrian Manea¹, Alexandru Schiopu^{1,3,4}

¹ Institute of Cellular Biology and Pathology “Nicolae Simionescu” of the Romanian Academy

² Department of Pathophysiology, University of Medicine, Pharmacy, Sciences and Technology of Targu Mures

³ Department of Translational Medicine, Lund University

⁴ Department of Internal Medicine, Skåne University Hospital

Background. The dysregulation of specific histone methylation-dependent epigenetic mechanisms leading to persistent transcriptional alterations, phenotypic switching, and trained immunity has been implicated in the etiology of cardiovascular disease. The SET7 and MLL1 histone methyltransferases induce histone methylation signatures associated with long-lasting transcriptional responses and monocyte hyperactivity. So far, the triggers of SET7 and MLL1-dependent regulatory mechanisms in myocardial infarction (MI) have not been investigated. S100A8/A9 is a neutrophil mediator that is highly increased in acute MI and drives local and systemic inflammatory responses.

Objective. We aimed to elucidate the potential role of S100A8/A9 in mediating SET7 and MLL1 expression and histone methylation in experimental MI.

Methodology. Male C57BL/6J mice were subjected to permanent left coronary artery ligation to induce MI. Sham-operated and MI mice were treated with 30 mg/kg of the selective S100A8/A9 inhibitor ABR-238901 or its vehicle, administered i.p. for three days. CD45+ cells were isolated from blood and bone marrow with a positive selection kit. Gene and protein expression was assessed by real-time PCR and Western blot.

Results. We found significant increases in SET7 and MLL1 mRNA/protein levels in the ischemic left ventricle and in CD45+ leukocytes from blood and bone marrow of MI mice. Concomitantly, the histone methylation markers reflecting the activation of SET7 (H3K4me1) and MLL1 (H3K4me3) were significantly elevated in the bone marrow-derived CD45+ cells. These epigenetic alterations were efficiently prevented by S100A8/A9 blockade with ABR-238901.

Conclusion. S100A8/A9 induces epigenetic modifications associated with trained immunity in the ischemic myocardium and bone marrow leukocytes of mice with MI. Pharmacological targeting of S100A8/A9 could be a promising therapeutic strategy to prevent long-lasting epigenetic alterations leading to leukocyte hyperactivity and persistent inflammation in acute MI.

Funding sources. Work supported by Romanian Academy and Romania’s National Recovery and Resilience Plan, PNRR-III-C9-2022-I8, CF148/15.11.2022, Financing Contract no. 760061/23.05.2023.

Financial disclosures. None.

Therapeutic S100A8/A9 inhibition reduces NADPH oxidase expression and counteracts NLRP3 inflammasome priming and activation in the ischemic myocardium

Mihaela-Loredana Vlad¹, Razvan Gheorghita Mares², Gabriel Jakobsson^{3,4}, Simona-Adriana Manea¹, Alexandra-Gela Lazar¹, Mihai Bogdan Preda¹, Maya Simionescu¹, Alexandru Schiopu^{1,3,4}, Adrian Manea¹

¹ Institute of Cellular Biology and Pathology “Nicolae Simionescu” of the Romanian Academy

² Department of Pathophysiology, University of Medicine, Pharmacy, Sciences and Technology of Targu Mures

³ Department of Translational Medicine, Lund University

⁴ Department of Internal Medicine, Skåne University Hospital

Background. Enhanced formation of reactive oxygen species (ROS) and alterations in redox signaling have been implicated in the pathophysiology of myocardial infarction (MI). The NADPH oxidase (Nox) complex is an important source of ROS in the infarcted myocardium. The alarmin S100A8/A9, primarily secreted by activated neutrophils, amplifies acute myocardial inflammation in MI and has previously been shown to be a promising therapeutic target to improve cardiac function post-MI.

Objective. We aimed to elucidate the underlying mechanisms linking S100A8/A9 and the inflammatory response in MI.

Methodology. MI was induced by permanent left coronary artery ligation in C57BL/6J mice, followed by treatment with the S100A8/A9 inhibitor ABR-238901 (30 mg/kg) or PBS for 3 days. The *in-vivo* experiments were complemented with mechanistic studies on cultured human macrophages (Mac), important effectors of inflammation and repair in the ischemic myocardium.

Results. Compared to sham-operated animals, we detected significant increases in the Nox1/2/4 catalytic subunits and in the inducible nitric oxide synthase (NOS2) at mRNA and protein level in the left ventricle of MI mice. Concomitantly, elevated levels of phosphorylated p65/NFκB, active caspase-1, and cleaved IL-1β were found in the ischemic left ventricle. S100A8/A9 blockade prevented the up-regulation of Nox and NOS2, suppressed NF-κB activation and prevented the priming and activation of the NLRP3 inflammasome. *In-vitro*, S100A8/A9 treatment induced expression of Nox1/2/4/5 catalytic subtypes and NLRP3 priming in human Mac in a TLR4-dependent manner. These effects were efficiently counteracted by ABR-238901 and by pharmacological inhibition of Nox1/Nox4 activity.

Conclusion. In conclusion, we show that Nox upregulation is an important contributor to S100A8/A9-induced NLRP3 inflammasome priming and activation in acute MI, leading to IL-1β production and myocardial inflammation.

Funding sources. Work supported by Romanian Academy and Romania’s National Recovery and Resilience Plan, PNRR-III-C9-2022-I8, CF148/15.11.2022, Financing Contract no. 760061/23.05.2023.

Financial disclosures. None.

Manuka Honey and its Phenolic Component, Methyl Syringate, Mediate Neutrophil Inflammatory Behaviors for Potential Use as a Vascular Prosthetic Bioactive

Evan N. Main¹, James C. Huang¹ and Gary L. Bowlin¹

¹Department of Biomedical Engineering, University of Memphis, Memphis, TN, USA

Neutrophils use reactive oxygen species (ROS) activity and a specialized form of cell death named NETosis to kill and ensnare invading pathogens. A dysregulated response can be deleterious and lead to tissue damage and fibrosis at host-biomaterial interfaces. Manuka honey has demonstrated potent antibacterial properties and recently, anti-inflammatory potential.

It was hypothesized that applying potentially therapeutic compounds found in abundance within Manuka honey to pro-inflammatory neutrophils will reduce intracellular ROS activity and prevent NETosis. Additionally, concentrations of whole Manuka honey were also assayed as a comparison.

Using primary human neutrophils isolated from donor (n=5) peripheral blood, concentrations between 1 nM and 1 mM of each flavonoid, 10 μ M and 2mM of methyl syringate, and 0.1% v/v and 10% v/v Manuka honey were assayed for reductions in NETosis using Sytox orange extracellular DNA staining and reduction in intracellular ROS activity via standard dichloro-dihydro-fluorescein diacetate (DCFH-DA) oxidation assay.

Results were expressed as relative ROS activity or NET formation (percentage) compared to positive control (stimulated cells) with standard deviation (n=5). Data were pooled, tested for normality, and statistically compared to positive control using Wilcoxon rank sum tests.

Whole Manuka honey reduced NET levels in an increasing pattern across all concentrations up to 91%, but only reduced ROS activity by 36% in a narrow concentration range. Methyl syringate, however, reduced NET levels by up to 68% and ROS activity by 66% within a broad therapeutic window of 10-1700 μ M.

In summary, the data from this study demonstrate that methyl syringate and whole Manuka honey are potent modulators of NET formation. In comparison with whole Manuka honey, methyl syringate is much more effective at lowering intracellular ROS activity. The results of the methyl syringate study versus other Manuka honey components indicates that methyl syringate drives the anti-inflammatory capabilities of Manuka honey demonstrated in previous studies.

Atypical melanocytes contribute to murine aortic valve elastogenesis

Perony da Silva Nogueira¹, Daniel Chaparro², Sana Nasim², Joshua Hutcheson² and Lidia Kos¹

¹Department of Biological Sciences, Florida International University, Miami, FL, US.

²Department of Biomedical Engineering, Florida International University, Miami, FL, US.

The aortic valve (AoV) opens and closes over two billion times in an average lifetime, regulating the blood flow from the left ventricle to the rest of the body. The remarkable ability to withstand the repetitive loading and unloading derives from the extracellular components, such as elastin (Eln) and collagen, that are maintained by specialized valvular interstitial cells (VICs), a heterogeneous cellular population within aortic valve leaflets. Pigment producing cells are observed in murine aortic valves and cells with traditional melanocytic markers exist within the interstitium of human aortic valve leaflets; however, the role of melanocytes in AoV development and disease remains largely unexplored. Our recent publications suggest a potential link between pigment production and elastin patterning within the mouse AoV. To further explore the role of melanocytes in patterning the developing AoV, we performed spatial transcriptomic analysis on murine AoV leaflets at postnatal day 3, focusing on genes closely linked to Eln expression. As expected, genes associated with smooth muscle cells (*Acta2*) and those critical for elastic fiber assembly (*Fbln5*, *Lox*) showed strong correlations with Eln. We also observed moderate correlations with melanocytic markers such as *Tyrp1* and a weaker correlation with the endothelial marker VE-cadherin (*Cdh5*). Immunolabeling of common phenotypic markers following *in situ* hybridization to identify Eln expression within AoV leaflets revealed populations of smooth muscle cells, endothelial cells, and melanocytes responsible for Eln production. Lineage tracing experiments using Wnt1-Cre/ZsGreen and Nkx2.5-Cre/YFP mice revealed that AoV melanocytes originate from both neural crest and second heart field lineages, providing the first evidence of non-neural crest-derived melanocytes within the developing heart. Future studies will explore the mechanistic contributions of different melanocytic populations to elastic fiber formation in the AoV.

Enhanced Targeted Repair of Vascular Injury by Apoptotic-cell-Mimicking Nanovesicles Engineered with P-Selectin Binding Peptide

Ruixin Zhang, Yu Shi, Yueyue Yang, Junwu Liu, Quhan Cheng, Deling Kong, * Kai Wang*

Key Laboratory of Bioactive Materials for the Ministry of Education, College of Life Sciences, Nankai University, Tianjin 300071, China.

E-mail: kongdeling@nankai.edu.cn (D. Kong); 013053@nankai.edu.cn (K. Wang)

Background and Objective: Late-stage coronary heart disease can lead to coronary artery occlusion, and treatment options usually involve percutaneous coronary intervention (PCI) with vascular injury as an almost inevitable consequence. Modulating inflammation and promoting re-endothelialization are crucial for repairing vascular injury. Phagocytosis of apoptotic cells at the injury site represents an effective mechanism for attenuating inflammation and improving regeneration during the natural healing process. However, strategies for repairing vascular injury using biomaterials derived from apoptotic cells have not yet been developed.

Method and Results: In this study, apoptotic body-mimetic nanovesicles (ApoNVs) derived from rat adipose-derived mesenchymal stem cells (rASCs) were prepared using a one-step extrusion method. The obtained ApoNVs inherit the unique anti-inflammatory and pro-regenerative properties of the parental apoptotic rASCs (Apo-rASCs). ApoNVs induce polarization of M1 macrophages toward the M2 phenotype. Meanwhile, they promote the proliferation, migration, and tube formation of endothelial cells (ECs). Moreover, ApoNVs-induced repolarized macrophages inhibit the proliferation and migration of vascular smooth muscle cells (VSMCs), and upregulate the functional gene expression of VSMCs. After engineering ApoNVs with P-Selectin Binding Peptide (ApoNVs-PBP), ApoNVs-PBP effectively target activated ECs, suppressing inflammatory gene expression. In a rat wire-mediated femoral artery injury model, ApoNVs-PBP further demonstrates targeted therapeutic effects on vascular injury.

Conclusion: ApoNVs-PBP effectively repairs vascular injuries by modulating macrophage phenotype, enhancing endothelial cell angiogenesis, and reducing inflammation, thereby offering a novel therapeutic strategy based on apoptotic cell.

Funding: This work was financially supported by the National Natural Science Foundation of China (81921004)

Material properties and acute human blood exposure may predict limb graft occlusion in EVAR

Michael F. Wolf, Hannah Jeffers, Job van Helvoirt, and Simon Overeem

Background: Abdominal Aortic Aneurysm (AAA) is the most common cardiovascular disease affecting the aorta and is typically treated with EndoVascular Aortic Repair (EVAR). A prevalent post-EVAR complication is limb graft occlusion (LGO), where thrombi block one or more stent limbs, often leading to rehospitalization. Literature shows different LGO rates between commercial stent grafts, with Zenith Alpha® having significantly higher rates compared to Endurant® and Excluder® [1]. No fundamental research has linked clinical parameters (kinking, reduced diameters, etc.) to the increased risk of LGO with certain devices. Hence, this research investigated the thrombogenicity of the stent fabrics as an explanation.

Methods: The thrombogenicity of Zenith Alpha®, Endurant®, and Excluder® devices was assessed according to ISO 10993-4 and compared using published methods and a dynamic test tube model [2]. Fabrics of the three devices were put in contact with fresh human blood from four donors, at two heparin levels, for one hour. The blood was then quantitatively analyzed (platelet count (Plt), thrombin-anti-thrombin (TAT), beta-thromboglobulin (βTG), and SC5b9), and fabrics were visualized with digital photography and scanning electron microscopy (SEM). ANOVA and Tukey-Kramer HSD analyses were applied.

Results: Analysis showed Zenith Alpha® being the most thrombogenic, with high platelet activation and thrombus formation, followed by Endurant® with significantly lower TAT, and Excluder® being the least thrombogenic with lowest platelet activation and thrombus formation based on significantly lower TAT, βTG release, and higher Plt ($P < 0.05$). SEM visualized high clot formation on Zenith Alpha®, mainly red blood cell attachment on Endurant®, and almost no thrombus on Excluder®.

Conclusion: These results show that the thrombogenicity of the materials correlates with the clinical LGO rates. This suggests that the previously uninvestigated thrombogenicity of stent graft materials may be the underlying cause of thrombotic obstructions resulting in Limb Graft Occlusions.

Funding: Medtronic.

Disclosure: Research by Medtronic.

References:

[1] <https://pubmed.ncbi.nlm.nih.gov/34266764/>

[2] <https://onlinelibrary.wiley.com/doi/full/10.1002/jbm.b.34705>

On the Role and Mechanisms of Notch Signaling in Endothelial-to-Mesenchymal Transition in Cardiovascular Diseases

Daria Evensen^{1,2}, Luisa Marracino³, Francesca Fortini⁴, Francesco Vieceli Dalla Sega⁴, Paola Rizzo⁵, Anna Malashicheva⁶, Jarle Vaage^{1,2,7}

¹Division of Physiology, Department of Molecular Medicine, Institute of Basic Medical Sciences, University of Oslo, Oslo, Norway

²Department of Research and Development, Division of Emergencies and Critical Care, Oslo University Hospital, Oslo, Norway

³Department of Translational Medicine and Laboratory for Technologies of Advanced Therapies, University of Ferrara, Ferrara, Italy.

⁴Maria Cecilia Hospital, GVM Care & Research, Cotignola, Italy.

⁵Maria Cecilia Hospital, GVM Care & Research, Department of Translational Medicine and Laboratory for Technologies of Advanced Therapies, University of Ferrara, Ferrara, Italy.

⁶Laboratory of Regenerative Biomedicine, Institute of Cytology Russian Academy of Science, Saint Petersburg, Russia.

⁷Institute of Clinical medicine, University of Oslo, Oslo, Norway

Background. Endothelial-to-mesenchymal transition (EndMT) is an important process in cardiac development and tissue maintenance. EndMT is also involved in the development of various cardiovascular diseases such as aortic valve calcification. Notch signaling is suggested to play a key role in EndMT. However, the exact mechanisms are still unclear.

Objectives. 1. To investigate the mechanisms of Notch activation or inhibition of EndMT. 2. To study the mechanisms of Notch activation and inhibition in endothelial cells.

Methods. Notch signaling was activated via lentiviral transduction of either Notch intracellular domain (NICD) or by ligand JAGGED 1 (JAG1). Notch signaling was down-regulated by short-hairpin RNA against either CSL (CBF1, Suppressor of Hairless, Lag-1) or against JAG1. CSL is the transcription factor in the cell nucleus which is activated by NICD. CSL is crucial for most Notch functions. Luciferase reporter assay was applied to investigate the interactions between either NICD or JAG1 with CSL. Immunocytochemistry was applied to verify EndMT.

Results. NICD induced expression of NOTCH2 and NOTCH3 as well as the downstream genes HEY1 and SLUG, followed by EndMT. JAG1 alone did not induce EndMT, causing only slight expression of Notch components. Unlike JAG1, NICD dose-dependently increased activity of the CSL luciferase reporter. Downregulation of CSL inhibited Notch as well as EndMT.

Conclusions. It is possible to activate and inactivate EndMT in the endothelial cells by activation and inactivation of Notch pathway components. Different ways of activation and inactivation of Notch signaling in endothelial cells have different effects on downstream signaling.

MICROPATTERNED SURFACE INVESTIGATION TO REDUCE THE RISK OF THROMBUS FORMATION IN CARDIOVASCULAR DEVICES

Marta Bonora (1), Veronica Viola (1), Stjepan Perak (2), Markus Lunzer (2), Sonja Kopp (3), Michael Mühlberger (3), Barbara Messner (4), Francesco Moscato (1), (5), (6)

1. Center for Medical Physics and Biomedical Engineering, Medical University of Vienna, Austria; 2. UpNano GmbH, Austria; 3. PROFACOR GmbH, Austria; 4. Department of Cardiac Surgery, Medical University of Vienna, Austria; 5. Ludwig Boltzmann Institute for Cardiovascular Research, Austria; 6. Austrian Cluster for Tissue Regeneration, Austria;

Background, hypothesis, and objectives

Implantable cardiovascular devices succeed in treating cardiovascular diseases. However, hemocompatibility-related issues still occur, and anticoagulant treatments are necessary 1–2. The objective of this study is to develop, through innovative technologies, an improved hemocompatible surface modified via specific micropatterns aimed at reducing the adhesion of platelets. Design implementation of the interface between biological tissue and artificial material may be the key point to reduce the related adverse events.

Methods

4 microgeometries were designed based on three criteria: i) hydrophobicity ii) reduction of accessible area for platelets adhesion iii) improvement of the blood Wall Shear Stress (WSS) effects. Micropatterns were printed via 2-photon-polymerization 3d-printer (2PP, UpNano GmbH) and replicated via Nanoimprinting Lithography (NIL, Profactor GmbH). Surface wettability and platelet adhesion were evaluated via Water Contact Angle (WCA) and Static adhesion tests.

Results

Geometries were printed in 4 dimensions: Extra-Small (1-3 μ m), Small (3-9 μ m), Medium (6–18 μ m), and Large (12-36 μ m). Wettability tests, compared to a flat surface showed higher hydrophobicity for Large and Medium sizes of all structures (WCA 120-130°) compared to Extra-Small and Small structures (WCA 100-110°). All the WCAs resulted in at least hydrophobic behaviour contrary to the flat surface (WCA 69°). Secondly, micropatterned surfaces were incubated with human platelets in physiological conditions, and compared to flat surfaces. All the micropatterned surfaces (16 samples) showed a significant reduction in platelet adhesion (between 20 – 50 %).

Conclusions

Micropatterned surfaces are highly hydrophobic, this feature may play an important role in platelet adhesion reduction through blood repulsion. This result agrees with platelet adhesion tests results. Platelet adhesion is favoured on flat surfaces. Micropatterned surface may be the solution to address this issue identified as first step of the thrombus formation.

Microfluidic experiments, aiming to test microstructure efficiency for a range WSS magnitudes experienced in blood pumps, are currently ongoing.

References

1. Li, Y. et al. A new way to evaluate thrombotic risk in failure heart and ventricular assist devices. *Med Nov Technol Devices* 16, (2022).
2. Vitale, N. et al. Left Ventricular Assist Device Thrombosis: Combined Approach by Echocardiography and Logfiles Review for Diagnosis and Management. *Braz J Cardiovasc Surg* 37, (2022).

Acknowledgements

The authors acknowledge funding from the OptiFlow 3D project (FFG FO999891239).

Surface modification with Adiponectin improves reendothelialization of synthetic small-diameter vascular grafts

Anna-Maria Schmitt^{1,2,3}, Sabrina Rohringer^{1,2,3}, Christian Grasl^{2,4}, Bruno K. Podesser^{1,2,3}, Helga Bergmeister^{1,2,3}

¹ Center for Biomedical Research and Translational Surgery, Medical University of Vienna, Vienna, Austria

² Ludwig Boltzmann Institute for Cardiovascular Research, Vienna, Austria

³ Austrian Cluster for Tissue Regeneration, Vienna, Austria

⁴ Center for Medical Physics and Biomedical Engineering, Medical University of Vienna, Vienna, Austria

Over the past decades significant progress have been made in the field of small-diameter vascular graft (SDVG) research. However, clinical issues related to insufficient endothelialization, thrombosis formation and intimal hyperplasia remain unsolved.

The aim of this project was to modify synthetic grafts to improve early reendothelialization. Electrospun thermoplastic polyurethane (TPU) grafts were loaded with Adiponectin via Click Chemistry.

By stimulating the phosphorylation of AMPK, Adiponectin downregulates the expression of pro-inflammatory mediators such as NFκB and TNF-α. It was shown that Adiponectin improves endothelial cell function by upregulating the expression of nitric oxide (NO) to prevent atherosclerosis development, while stimulating COX-2 and thereby reducing apoptosis. Adiponectin further protects intimal hyperplasia by suppressing smooth muscle cell (SMC) proliferation through inhibition of growth factors, making it an interesting therapeutic target in vascular graft research.

The influence of Adiponectin on interactions between cells and the synthetic material was tested *in vitro* by seeding rat aortic endothelial cells (RaOECs) onto the modified surface. Cell viability assessments revealed EC proliferation over time and proved cytocompatibility of Adiponectin-loaded TPU grafts. Immunofluorescence staining demonstrated successful reendothelialization that was further confirmed by electron microscopy. Gene expression analyzed by qPCR revealed no significant changes in cellular expression patterns.

The results of this study demonstrated that Adiponectin promotes the development of an intact endothelium by improving cell-graft interaction. We hypothesize that immobilization of Adiponectin on synthetic SDVGs could have a great impact on early healing processes.

This project was funded by the Ludwig Boltzmann Institute of Cardiovascular Research.

Salvianolic Acid A (SA) fosters lymphangiogenesis against myocardial ischemic reperfusion injury

Zhuo LIU¹, Ruizhi LI², Wenxin HUANG¹, Hong ZENG¹, Ying QU¹, Tonghe ZHU², Chen XU*

1 Department of Physiology and Pathophysiology, Shanghai Key Laboratory of Bioactive Small Molecules, School of Basic Medical Sciences, Fudan University, 138 Xueyuan Rd., Shanghai 200032, P.R. China

2 Multidisciplinary Centre for Advanced Materials, Institute for Frontier Medical Technology, School of Chemistry and Chemical Engineering, Shanghai University of Engineering Science, 333 Longteng Rd., Shanghai 201620, P.R. China

*Corresponding author: Chen XU (xuchenfd@fudan.edu.cn)

Abstract: All abstracts should include: (1) a brief introduction, (2) relevant information on methods, (3) summarized results, and (4) conclusion drawn from the results.

Ischemic heart disease is one of the leading causes of death globally. The lymphatic system plays a key role in regulating tissue fluid homeostasis, lipid transport, and immune surveillance. In response to tissue damage, lymphangiogenesis could engender favorable microenvironment which highlights its tremendous potential as therapeutic strategies to improve MI prognosis. Salvianolic acid A (SA) is an extraction from traditional Chinese medicinal herb *Salvia miltiorrhiza*. Previously we developed degradable polyurethane elastomers with SA as a chain extender [1]. Based on it, we employed 3D printing and near-field direct writing technology to designed a novel cardiac patch (PEUU/SA/3D/Near-field, PD-DN), which resembling cardiac extracellular matrix in porosity, orientation and spatial arrangement. Featuring the biomimetic microstructure and persistent release of SA, PD-DN serves as a versatile scaffold for cell adhesion and climbing. In I/R injury SD rat models, we observed that PD-DN patch effectively promoted lymphangiogenesis, alleviates cardiac remodeling, preserved heart strain and remarkably ameliorated heart systolic function. Mechanistically, we predicted the potential targets of SA based on PharmMapper and SwissTargetPrediction databases. Among the top 10 core targets, SA competitively binds to the ATP binding pocket of epithelial growth factor receptor (EGFR), which could directly block the activation of tyrosine kinases and their downstream signaling. Utilizing single-cell RNA sequencing, we identified a unique subgroup of fibroblasts activated by SA in early-stage repair.

TGFB2 and Colla12a secreted from these fibroblasts activated TGFBRs and integrin pathways on lymphatic endothelial cell to foster early-stage lymphangiogenesis. Collectively, PD-DN sheds light on a feasible way to exploit therapeutic effects of SA while remedy its deficiency of non-specific targeting and short half-life period, which provides an enlightening prospect of bioengineering and clinical practice in ischemic heart diseases.

Keywords: Salvianolic Acid A, Tissue Engineering, Lymphangiogenesis, Ischemia/Reperfusion Injury

Acknowledgement: This work is supported by Innovative research team of high-level local universities in Shanghai and a key laboratory program of the Education Commission of Shanghai Municipality (ZDSYS14005); National Key Research and Development Program of China (2023YFC2412400, 2023YFC2412403); and Technology Support Project of Science and Technology Commission of Shanghai Municipality (22S31904700).

Reference:

[1] Liu Y, Wang L, Liu Z, Kang Y, Chen T, Xu C, Zhu T. Durable Immunomodulatory Nanofiber Niche for the Functional Remodeling of Cardiovascular Tissue. ACS Nano. 2024 Jan 9;18(1):951-971.

Development of a new protective solution for vein grafts in bypass surgery

Tobias Braun¹ und Theodor Fischlein¹
Maria Kokozydou² und Gundula Schulze-Tanzil²

¹ Department of Cardiac Surgery, Klinikum Nürnberg, Paracelsus Medical University, Nuremberg, Germany

² Institute for Anatomy and Cell Biology, Paracelsus Medical University, Nuremberg, Germany

Clinical Problem

Bypass operations are often the only treatment option for atherosclerosis [1]. However, saphenous vein grafts (SVGs) frequently fail shortly after surgery. SVGs are stored in preservative solutions, which can lead to functional decline due to the type of preservation solution used [2, 3], surgical trauma, and other factors. Research has shown that saline, the most common solution, is inadequate for storing blood vessels, causing partial detachment of the endothelium [2]. This damage makes the graft more prone to platelet adhesion, macrophage infiltration, plaque development, and intimal hyperplasia, resulting in graft failure [5].

Objective

The aim is to develop a preservation solution that protects endothelial cells (ECs) from cellular stress during bypass grafting. In our previous study, we compared several commonly used solutions. Our goal is to further optimize these solutions to protect endothelial integrity better than standard solutions.

Material and Methods

Remnants of SVGs were used for (immuno-)histological staining to detect endothelial detachment or to isolate ECs for in vitro experiments (e.g., cell viability assays, metabolic activity measurements, migration capability tests, and activity assessments of endothelium-specific genes). Additionally, ex vivo experiments with the vein remnants are planned.

Results

Our findings indicate that full electrolyte solutions with an extracellular composition are more suitable for preserving SVGs compared to saline solution (that showed the overall lowest protection) or solutions with a low sodium content. Additionally, metabolizable buffer substances like lactate show potential in maintaining endothelial integrity by serving as an energy source and increasing migration capabilities.

Conclusions

The research aims to create a perfusion solution that meets the stringent requirements of cardiac and vascular surgery, improving long-term graft patency and patient outcomes. Such a solution could significantly reduce the failure rate of SVGs and enhance the success of bypass operations.

Funding Sources

This project is funded by PMU University funding.

[1] R.N. Singh, J.A. Sosa, G.E. Green, Internal mammary artery versus saphenous vein graft. Comparative performance in patients with combined revascularisation, *Br Heart J* 50(1) (1983) 48-58.

[2] D.R. Weiss, G. Juchem, M. Eblenkamp, B.M. Kemkes, B. Gansera, M. Geier, S. Nees, Search for optimized conditions for sealing and storage of bypass vessels: influence of preservation solution and filling pressure on the degree of endothelialization, *Int J Clin Exp Med* 3(1) (2010) 10-27.

[3] H.R. Zerkowski, M. Knocks, M.A. Konerding, N. Doetsch, G. Roth, K. Hakim, J.C. Reidemeister, Endothelial damage of the venous graft in CABG. Influence of solutions used for storage and rinsing on endothelial function, *Eur J Cardiothorac Surg* 7(7) (1993) 376-82.

[4] Jeffrey V. Manchio, BS, Junyan Gu, MD, Linda Romar, RN, James Brown, MD, James Gammie, MD, Richard N. Pierson III, MD, Bartley Griffith, MD, and Robert S. Poston, MD
Disruption of Graft Endothelium Correlates With Early Failure After Off-Pump Coronary Artery Bypass Surgery; Division of Cardiac Surgery, University of Maryland School of Medicine, Baltimore, Maryland

[5] Christopher D. Owens, Warren J. Gasper, Amreen S. Rahman, Michael S. Conte; Vein graft failure; *Journal of Vascular Surgery*; Volume 61, Issue 1; 2015; Pages 203-216; ISSN 0741-5214, <https://doi.org/10.1016/j.jvs.2013.08.019>.

In-vivo Evaluation of Biomimetic Vascular Grafts with Artery-Tuned Mechanical Properties

Kaspars Maleckis¹ and Jason MacTaggart²¹Department of Biomechanics, University of Nebraska Omaha, Omaha, Nebraska 68182, United States²Department of Surgery, University of Nebraska Medical Center, Omaha, Nebraska 68198, United States

Native arteries exhibit complex mechanical characteristics, including low stiffness, nonlinearity, and anisotropy, which vary substantially to meet local physiological demands. Synthetic vascular grafts currently lack these mechanical features, which can lead to suboptimal physiology, decreased functional remodeling, and increased failure rates. Despite ongoing efforts to develop mechanically biomimetic synthetic vascular grafts, existing biomaterials and manufacturing methods have not yet succeeded in achieving artery-tuned nonlinear and anisotropic mechanical properties.

This work aimed to develop and refine a manufacturing method that enables tuning stiffness, nonlinearity, and anisotropy of synthetic vascular grafts to specific arteries and evaluate the effects of mechanical biomimicry on graft performance. Electrospun Nanofibrillar Grafts (ENGs) were manufactured from a mixture of biomedical-grade elastomeric polyurethanes. ENG mechanical parameters were controlled by adjusting graft wall thickness, polymer ratio, and fiber alignment. ENG prototypes were then compared to commercial devices via benchtop assessments and preliminary in-vivo feasibility studies in domestic swine models for peripheral arterial and aortic repairs.

The developed synthetic graft manufacturing approach allows the control of ENG stiffness, nonlinearity, and anisotropy in the range of native artery properties. Benchtop studies demonstrate that the ENG prototypes have appropriate surgical properties and cytocompatibility. In-vivo studies reveal the potential benefits of mechanically tuned ENGs over commercial grafts in peripheral arterial and aortic repairs, evidenced by improved endothelialization, smooth muscle cell incorporation, and maintenance of biomimetic mechanical properties for up to 16 weeks.

The developed ENG manufacturing technology enables achieving a wide range of arterial mechanical characteristics that can be independently tuned to specific arteries. While further development and testing is warranted, this technology holds promise in developing synthetic vascular grafts with optimized biomechanics and improved clinical outcomes.

This research was partially supported by NIH R01HL147128, NIH P20GM152301, NIH R61HL173890, and the University of Nebraska Collaboration Initiative grant #27311.

Reinforced Biotubes as Readily Available and Regenerative Vascular Grafts

Quhan Cheng, Dengke Zhi, Kai Wang, Deling Kong*

Key Laboratory of Bioactive Materials for the Ministry of Education, College of Life Sciences,
Nankai University, Tianjin 300071, China

E-mail: kongdeling@nankai.edu.cn

Background and Objective: There is a great clinical demand for vascular substitutes that can achieve vascular regeneration in vascular replacement or bypass for cardiovascular disease, and arteriovenous graft (AVG) for hemodialysis. Decellularized extracellular matrix (dECM) is a promising biomaterial for tissue regeneration. Herein, we sought to develop a decellularized vascular substitute with the capability of recellularization and promoting vascular regeneration.

Method and Results: Decellularized and polymer fiber skeleton (PS) reinforced biotubes (dPB) was constructed by subcutaneously embedding the PS and followed decellularization. The microstructure of dPB was assessed by the scanning electron microscope, which showed that the dECM in dPB was integrated with PS and had a loose and porous microstructure, enabling dPB excellent performance of recellularization after implantation. Proteomics analysis and *in vitro* cell assay showed that dPB could promote the macrophage polarization to anti-inflammatory phenotype, and further promoted vascular regeneration. Long-term *in vivo* evaluation of dPB were performed in different animal model. dPB was demonstrated the capability of promoting vascular endothelial and smooth muscle regeneration, recovering physiological functions, anti-infection and resistance to needle puncture in the models of canine carotid artery replacement, pig coronary artery bypass graft (CABG), and canine AVG. The initial safety and efficacy were validated by using as AVG in a first-in-man clinical trial.

Conclusion: This work offers an off-the-shelf and pro-regenerative dECM vascular graft for future possible wide use in range of vascular applications for patients with no available autologous vascular substitutes.

Funding: This study was funded by the National Natural Science Foundation of China (NSFC) projects (81921004 and 82172106).

Complete Transformation of Bioresorbable Synthetic Vascular Grafts in the Common Carotid Artery

Anthony D'Amato¹, Cole Latvis¹, Nathaniel Wright¹, Ying Chen¹, Edith Tzeng², Yadong Wang¹

1: Nancy E. and Peter C. Meinig School of Biomedical Engineering, Cornell University, Ithaca, USA

2: Department of Surgery, Univ. of Pittsburgh, Pittsburgh, PA, USA

When arteries occlude, rerouting blood flow with a vascular graft can save limbs and lives. Autologous grafts are the preferred conduit for replacing small-diameter vessels. However, not all patients have suitable donor vessels, and the wound healing complications associated with the harvest can be severe. Synthetic grafts can be used but the foreign material does not degrade or integrate with the host, leading to graft failure and risk of infection. These grafts function best in large caliber arterial reconstruction but perform poorly in smaller arteries such as in the coronary and below the knee arteries. We believe that by improving the materials used in vascular grafts we can overhaul their performance and expand their use in the clinic. We are approaching this challenge by creating acellular, synthetic vascular grafts using a novel elastomer. This material is processed to create highly porous grafts that have been implanted in both rat and sheep common carotid interposition models. We have successfully demonstrated complete transformation of these synthetic grafts to autologous tissue in both animal models – with the presence of elastin in the newly regenerated artery – a feat that to our knowledge has never been achieved in large animals before. Successful translation of this technology stands to benefit millions of patients every year that require below the knee bypass, coronary artery bypass, and arteriovenous graft placement, to name a few. This work is ongoing and is supported by NIH Grant #R01HL159427. This technology is also currently being commercialized through a startup out of Cornell University.

aXess: Progress update on a restorative hemodialysis access vascular conduit

Martijn Cox¹, Frederick J. Schoen², Renu Virmani³, Paulo Neves¹, An de Vriese⁴

¹Xeltis BV, Eindhoven, The Netherlands

²Brigham & Women's Hospital and Harvard Medical School, Boston, USA

³CVPPath Institute Inc, Gaithersburg, USA

⁴Department of Nephrology and Infectious Diseases, AZ-Sint Jan, Brugge, Bruges, Belgium.

Current options for hemodialysis access have significant limitations for patients, who often face a cycle of complications and reinterventions. The Xeltis aXess hemodialysis access vascular conduit is composed of an absorbable polymer supported by an intramural nitinol microskeleton. It is designed to enable cell infiltration and new tissue formation to replace absorbed polymer, resulting in functional endogenous tissue restoration (ETR) and efficacious and safe repetitive puncturability. Ongoing preclinical and clinical studies suggest this approach may combine benefits of an off-the-shelf synthetic approach with the potential to harness the body's innate healing capacity.

A GLP preclinical study was performed using a sheep carotid-to-jugular-vein shunt model comparing 16cm long, 6mm inner diameter aXess conduits (1-, 3- and 6-months f-up) with size-matched ePTFE controls (6-months f-up). 24-months f-up on 2 additional aXess sheep is pending and will be available shortly (17 sheep total).

Primary patency at 6 months was 50% (3/6) for aXess grafts compared to 0% (0/3) for ePTFE graft controls. aXess conduits showed expected levels – mild to moderate - of host macrophage-mediated inflammation and polymer resorption starting from the abluminal aspect. Over time the bulk polymer separated into fragments that either replaced or encapsulated by new collagenous tissue, both associated with a reduced inflammatory response once the process locally completes. The remodeling response of grafts is consistent with other ETR applications of this polymer system.

Based on earlier preclinical studies, regulatory approval was obtained in Europe for a First-In-Human clinical study which has enrolled 20 patients, with 12 months follow-up completed for full cohort. Based on favorable safety data and patency rates, an EU pivotal clinical study was started and is now enrolling patients.

These studies demonstrate the potential of restorative polymer-based approaches to hemodialysis in humans, which may offer benefits compared to currently available options.

Financial disclosures: Cox and Neves are employed by Xeltis. Schoen is a paid consultant and Scientific Advisory Board member to Xeltis. Virmani is a paid consultant to Xeltis.

Inhibition of mitochondrial complexes I and IV reduces vascular calcification.

Andrea Gorgels, Tabea Koeneke, Marina Heuschkel, Claudia Goettsch

Department of Internal Medicine I – Cardiology, Medical Faculty, RWTH Aachen University, 52074 Aachen, Germany

Cardiovascular calcification (CV) is a serious global health issue. It independently increases the risk of CV disease and death with no effective treatments currently available. Factors contributing to vascular calcification include inflammation, endoplasmic reticulum stress, and metabolic imbalances in vascular smooth muscle cells (SMCs). These metabolic imbalances can cause or result from changes in mitochondrial function. Our recent research has demonstrated alterations in mitochondrial function in calcifying SMCs. The five mitochondrial complexes are key to mitochondrial activity. Therefore, we analyzed the different mitochondrial complexes, hypothesizing that their activities contribute differently to vascular calcification.

SMCs were calcified in osteogenic media (OM). We assessed the function of individual mitochondrial complexes using the Seahorse approach to investigate the metabolic impact of the calcifying environment. Through mitochondrial efflux analysis, we observed that calcifying SMCs displayed increased complex I oxygen consumption rate (OCR) (+39%, $p=0.025$) and complex IV OCR (+47%, $p=0.011$). Inhibiting complex I with rotenone and complex IV with potassium azide decreased matrix mineralization in calcifying SMCs dose-dependent (Rotenone -80%, $p=0.002$; Potassium azide -65%, $p<0.001$). Interestingly, complex II activation with succinate increased matrix mineralization (1.56-fold ($p=0.010$), which was prevented by rotenone, suggesting that complex II plays a minor role. Activating complex IV with TMPD did not affect matrix mineralization. Rotenone and potassium azide did not change ALPL mRNA expression and tissue non-specific alkaline phosphatase activity. However, rotenone and potassium azide reduced the extracellular pH without affecting the intracellular pH, suggesting that extracellular acidification may impact matrix mineralization. Subsequent analysis of extracellular lactate showed no changes.

In conclusion, complexes I and IV may contribute to vascular calcification, as blocking these complexes lowers SMC mineralization. This decrease in mineralization does not follow the traditional pathway associated with reduced ALPL mRNA expression and activity. The role of extracellular acidification in this mechanism requires further investigation.

This work was supported by the German Research Foundation (DFG SFB/TRR219-C02 to CG) and by the Excellence Strategy of the Federal Government and the Länder ERS RWTH Aachen (RWTH Start-Up StUpPD_448-23 to AG)

Title: Exploring the Role of Caveolin-1 in the Calcification Paradox: Cellular Insights into Divergent Mineralization Pathways in Vascular and Bone Tissues

Katherine Kaiser, Amirala Bakhshian Nik, Patrick Sun, Bohdan B. Khomtchouk, Joshua Hutcheson
Department of Biomedical Engineering, Florida International University, Miami, FL, United States

Though vascular smooth muscle cells (VSMCs) adopt an osteogenic phenotype during pathological vascular calcification, clinical studies note an inverse correlation between bone mineral density and arterial mineral—a phenomenon known as the calcification paradox. Both mineralization processes are mediated by extracellular vesicles (EVs) that sequester calcium and phosphate. Previous studies associate calcification paradox with an imbalanced phosphate/calcium content due to bone loss and/or increased systemic inflammation. However, modulation of caveolin-1 (CAV1), a membrane scaffolding protein essential to calcifying EV formation in the vasculature, also results in inverse effects on mineralization in bone and vascular tissues. Moreover, across two independent epidemiological cohorts, CAV1 genetic variation demonstrates inverse relationships between bone mineral density (BMD) and coronary artery calcification (CAC). These findings indicate that the calcification paradox may not be fully attributed to the systemic distribution of mineral components or inflammation. To further investigate this divergent role of CAV1 in mineralization, CAV1 expression was knocked down by siRNA in primary human coronary artery VSMC and human osteoblast (HOB) cell cultures, grown in both control and osteogenic media. Methyl β -cyclodextrin (M β CD) and a calpain inhibitor were used to disrupt and stabilize the caveolar domains, respectively, in VSMCs and HOBs. At the study endpoint, calcification was measured via Alizarin Red S staining. *In vitro* siRNA knockdown of CAV1 abrogated calcification production in VSMCs ($p < 0.05$), while having no effect on osteoblast mineralization. M β CD-mediated caveolae disruption led to a 3-fold increase of calcification in VSMCs treated with osteogenic media ($p < 0.05$) but hindered osteoblast mineralization ($p < 0.01$). Conversely, stabilizing caveolae by calpain inhibition prevented VSMC calcification ($p < 0.05$) but had no significant effect on osteoblast mineralization. Our data indicate fundamental cellular-level differences in physiological and pathophysiological mineralization mediated by CAV1 dynamics in VSMC and HOB cultures.

Discovering the Effects of Cancer Cells on Lymphatic/Blood Vessel MispatterningArinola O. Lampejo¹, Walter L. Murfee¹¹J. Crayton Pruitt Department of Biomedical Engineering, University of Florida, Gainesville, FL

The tumor microenvironment is known to induce microvascular remodeling, including both angiogenesis and lymphangiogenesis. Motivated by the challenge to recreate the multi-cell/system complexity of the tumor microenvironment, our laboratory recently developed an *ex vivo* biomimetic model that integrates tumor spheroids and cultured rat mesentery tissues. The model enables real-time observation of cancer cells, blood vessels, and lymphatic vessels and its application led to the discovery of tumor spheroid-associated lymphatic/blood vessel mispatterning. To validate the patho-physiological relevance of our model, the objective of this study was to determine the effects of tumor spheroids on microvascular remodeling post transplantation in analogous *in vivo* studies. H1299 human lung carcinoma cell lines were labeled with DiI and suspended in a hanging drop culture system consisting of MEM supplemented with 10% serum for 4 days to form fluorescent spheroids. These spheroids were then seeded onto exposed rat mesentery tissues. Tissues were then harvested 3- and 5-days post seeding to match *ex vivo* time points and labeled with BSI-lectin, PECAM, and/or LYVE-1. The spheroid group was compared to sham (no spheroids) and unstimulated tissues. On day 3 and day 5, clusters of DiI cancer cells were observed. A distribution of individual cells across the tissues suggested migration from their initial spheroid configurations. Evidence of cancer cell effects on angiogenesis and lymphangiogenesis was supported by increased blood capillary sprouting and density at both 3 and 5 days. 5-day spheroid tissues also displayed examples remodeled lymphatic vessels characterized by dramatic changes in morphology, apparent blebbing, segment specific increases in diameter, and examples of segment specific decreased LYVE-1 labeling. The results support the relevance of using our *ex vivo* model to guide *in vivo* investigation and offer a new view for understanding altered microvascular structure and function in a tumor microenvironment. Funding source: American Heart Association Grant 23TPA1142184.

Algorithm-Driven Optimization of Helical Vascular Grafts to Suppress Neointimal Hyperplasia in Arteriovenous Grafts

Boxin Liu, Deling Kong*, Kai Wang*

Key Laboratory of Bioactive Materials for the Ministry of Education, College of Life Sciences,
Nankai University, Tianjin 300071, China

Background and Objective: There is a high clinical demand for arteriovenous grafts (AVGs) that can inhibit intimal hyperplasia and maintain long-term patency in hemodialysis. Helical flow can enhance the hemodynamic performance of grafts, thereby suppressing intimal hyperplasia and thrombosis at the anastomotic site. However, the optimal helical structure design remains unachieved. Therefore, we aim to develop an efficient helical artificial blood vessel capable of inhibiting intimal hyperplasia and maintaining long-term patency by combining finite element methods and mathematical statistical algorithm.

Methods and Results: We constructed computational models of AVGs with various helical structures (including anastomotic angle, pitch, and ridge height). Using finite element analysis, we extracted key hemodynamic parameters from the anastomotic region of these helical AVGs. Mathematical statistical methods were then employed to analyze and optimize the hemodynamic parameters of each model, leading to the design of a helical blood vessel structure that provides optimal hemodynamic conditions. The optimized helical blood vessels were fabricated according to the design parameters using a combination of melt spinning and electrospinning. Mechanical characterization showed that the fabricated blood vessels met the requirements for artificial vascular graft transplantation, including burst pressure, suture strength, and mechanical strength. *In vivo* evaluation was conducted using a pig AVG model, which demonstrated that the optimized helical artificial blood vessels could induce helical flow *in vivo* and exhibit excellent capability to inhibit intimal hyperplasia, thereby maintaining long-term patency.

Conclusion: This study provides a new method and approach for the structural optimization of artificial blood vessels. The optimized helical artificial blood vessels, which effectively inhibit intimal hyperplasia and support long-term patency, hold promise for applications in various cardiovascular diseases requiring artificial vascular graft transplantation.

Funding: This study was funded by the National Natural Science Foundation of China (NSFC) projects (82172106).

Creating order out of chaos – Modelling the influence of chronic inflammation on restoring cardiac structural anisotropy

Marjolein J.M. ten Dam (m.j.m.t.dam@tue.nl)¹, Bart G.W. Groenen (b.g.w.groenen@tue.nl)¹, Anthal I.P.M. Smits (a.i.p.m.smits@tue.nl)¹, Carlijn V.C. Bouten (c.v.c.bouten@tue.nl)¹

¹*Technical University Eindhoven, Soft Tissue Engineering and Mechanobiology*

Heart attacks affect one in five men and one in six women, posing a significant health concern.¹ After myocardial infarction (MI), cardiac function declines due to the loss of cardiomyocytes, which is later replaced by disorganized scar tissue. Recent studies from our group suggest that restoring tissue to its natural, anisotropic organization significantly improves cardiac function. However, it remains unclear whether tissue anisotropy can be restored in the presence of chronic inflammation - typical for infarcted myocardium. This study will be devoted to unravelling the impact of chronic inflammation on restoring anisotropy. We hypothesize that chronic inflammation will hamper tissue anisotropy restoration due to adverse tissue remodelling, as it dysregulates the ECM turnover and cellular activation.

To adequately mimic tissue organization, we utilize a mini-tissue model in which cells are seeded in between two Velcro constraints for anisotropic organization, and four constraints for chaotic organization. Cardiac fibroblasts and cardiomyocytes are suspended in a 30:70 ratio within a gel composed of collagen type I and Matrigel. Tissue contractility is enhanced through electrical pacing. To study remodeling, the four-constraint tissue is cut from two opposing Velcro constraints and observed for one week. During this period, tissues are exposed to various cocktails of pro- and anti-inflammatory cytokines to mimic chronic post-MI inflammation. At different time points, cellular and collagen alignment will be analyzed using our in-house developed CNA-OG488 probe and cell-specific markers. Additionally, cell proliferation and activation will be assessed using qPCR, immunofluorescence, and DNA content measurement with the dsDNA BR assay kit. Hydroxyproline will be quantified with a Chloramin-T assay. Mechanical properties and rupture behavior will be evaluated through uniaxial tensile testing. Finally, cardiac contractility will be assessed using a calcium transient assay.

This model allows for systematic dissection of the influence of inflammatory cytokines on structural remodeling and, consequently, cardiac function.

This project is funded by the ERC advanced grant RE-ALIGN.

¹American Heart Association, Women vs Men Heart Attack

Lymphatic and Tenascin C Crosstalk in Left Ventricular Hypertrophy-Induced Cardiac Remodeling

Lukas Weber^{1*}, Simge Baydar^{1*}, Zsombor Ocskay^{3*}, Anna Stampfer¹, Peter Pokreisz¹, Karin Zins², Dietmar Abraham², Bruno K. Podesser¹, Zoltan Jakus^{3*}, Attila Kiss^{1*}

¹Ludwig Boltzmann Institute for Cardiovascular Research at the Center for Biomedical Research and Translational Surgery, Medical University of Vienna, Austria

²Center for Anatomy and Cell Biology, Division of Cell and Developmental Biology, Medical University of Vienna, Austria

³Department of Physiology, Semmelweis University School of Medicine, Budapest, Hungary

*Contributed equally to this work.

Introduction: Boosting lymphangiogenesis has proven to alleviate inflammation, fibrosis and myocardial edema after myocardial infarction (MI) and induce overexpression of Tenascin-C (TNC). TNC is a matricellular protein that negatively regulates lymphangiogenesis post-MI, contributing to impaired repair and remodeling process. Nevertheless, the dynamic changes in lymphangiogenesis in pressure-overload induced left ventricular hypertrophy (LVH) and its crosstalk with TNC are poorly understood.

Methods: LVH was induced by transverse aortic constriction (TAC) in adult male C57BL/6 mice, allocated to groups of sham, 1-week and 6-weeks post-TAC. Hearts were processed for histology of Masson's trichrome and immunostainings for lymphatic markers (Podoplanin and LYVE1) and TNC. Alterations of lymphatic vessels were quantified in perivascular areas and total heart tissue. In addition, lymphatic endothelial cells (LEC) were subjected to 48h low (1-3%) or high (18-21%) elongation using FlexCell to mimic stretch overload. The expression of NF- κ B and fibrotic markers (Collagen I and III) were assessed by western blot and RT-qPCR. Scratch-wound healing and XTT cell viability assays were performed on LEC after treatment with recombinant human TNC for 24h (1 μ g/ml or 5 μ g/ml).

Results: Region-specific alterations of lymphatic vessels in heart tissue after pressure overload were detected. In addition, 1-week post-TAC, a regression of lymphatic structures was observed exclusively in perivascular areas, along with increased perivascular fibrosis and TNC expression. High stretch conditions in LEC showed increased expression of inflammatory and fibrotic markers, suggesting lymphatic remodelling. Scratch-wound healing and XTT assays revealed prolonged wound healing and a tendency towards reduced viability in LEC after TNC treatment.

Conclusion: Our study demonstrates dynamic spatio-temporal changes of lymphatic vessels in LVH due to pressure overload. Moreover, TNC appears to be a crucial player in modulating lymphatic vessel growth in LVH. Thus, inhibiting TNC and/or boosting lymphangiogenesis may be a potential novel therapeutic strategy to alleviate fibrosis and remodeling in LVH.

Carnosine-copper chelator-modified small-diameter vascular grafts for the promotion of anticoagulation and endothelial regeneration

Shaojin Lei, Shufang Wang*

Key Laboratory of Bioactive Materials, Ministry of Education, College of Life Sciences, Nankai University, Weijin Road 94, Tianjin 300071, P. R. China

Corresponding author Email: wangshufang@nankai.edu.cn

Abstract: Thrombosis and poor endothelialization of small-diameter vascular grafts (SDVGs) remain the leading cause of frequent failure, despite current tissue-engineered vascular grafts providing a feasible strategy in clinical treatment strategies for the small-diameter (<6 mm) blood vessels in management of cardiovascular disease. Copper ions are normally used to achieve the purpose of antithrombotic because they can produce nitric oxide (NO) with the catalysis of NO donors. However, copper ions also participate in the Fenton reaction to produce reactive oxygen species (ROS), which would adversely affect vascular graft remodeling. Here carnosine, an ion-chelating agent with anti-inflammatory and antioxidant properties, was used to chelate with copper ions to synthesize carnosine-Cu (II) chelate which showed effective activity of antithrombotic, anti-inflammatory, and antioxidant. Besides, our research further demonstrated that copper ions still retain the catalytic activities for producing NO after chelating with carnosine accompanied by the production of lower levels of ROS (the average fluorescence intensity of ROS decreased from 79.69 to 7.13). Furthermore, the grafts achieved excellent patency with satisfactory endothelialization (the endothelial coverage reached 80.64%) and significantly improved deposition of collagen and glycosaminoglycan for up to 12 weeks. In general, Car-Cu (II) modified SDVGs provided a facile approach to addressing the problems of thrombus and poor endothelialization in SDVGs for the clinic, which is significant in cardiovascular regenerative medicine.

Funding sources: This study was supported by Tianjin Research Program (22JCZXJC00080, S22ZDF291), the National Key Research Development Program of China (2020YFA0803701).

Simulated microvasculature in thick 3D bioprinted collagen-based matrices promoted in active perfusion bioreactor

Denisa Kanokova¹, Jana Matejkova¹, Monika Supova², Jaroslav Chlupac³, Roman Matejka¹

¹ Department of Biomedical Technology, Faculty of Biomedical Engineering, Czech Technical University in Prague, Kladno, Czech Republic

² Department of Composites and Carbon Materials, Institute of Rock Structure and Mechanics, Czech Academy of Sciences, Prague, Czech Republic

³ Transplantation Surgery Department, Institute for Clinical and Experimental Medicine (IKEM), Prague, Czech Republic

The 3D bioprinting of high-concentrated collagen bioinks is a promising technology for tissue engineering and regenerative medicine involving creating new tissue models or replacements. Due to its natural abundance in the extracellular matrix of many tissues and its biocompatibility, collagen is a widely used biomaterial for bioprinting. High-concentrated collagen hydrogels have shown great potential due to their favorable structural and biomechanical properties. However, achieving high cell proliferation rates within these hydrogels remains challenging, especially for thick (> 1.5 mm) samples, due to reduced diffusion and exchange of nutrients and metabolites. In this study, we aimed to prepare 2 mm thick collagen matrices with incorporated stromal/stem cells with maximum viability and their ability to differentiate into smooth muscle cells and endothelial-like cells. The collagen bioinks were based on porcine collagen isolated from porcine skins. We have optimized the procedure for printing at densities of 20, 30, and 50 mg/mL with incorporated porcine adipose-derived stromal cells and Wharton's jelly stem cells. We then used several approaches in the cultivation and creation of microvasculature. Static cultivation of the samples showed minimal viability only in border layers, even for samples 1 mm thick. We then utilized controlled active perfusion with freely floating samples, maintaining whole-volume wetting and culture media exchange. This resulted in viable cells in 1 to 1.5 mm thick samples. Punctured and print-patterned microchannels were then incorporated. This approach increased the overall diffusion area, promoting cell viability in thick samples. Downsizing reduces mechanical stability and requires precise preparation of the microchannels because they are prone to collapse. However, combining microchannels in collagen matrices with active perfusion in a bioreactor showed promising results in thick sample preparation. In addition, cells were able to remodel the substrate by differentiating into SMC-like structures.

This research was funded by the Ministry of Health of the Czech Republic grant No. NW24-08-00064 and NW24J-02-00061.

Supporting Autologous Internal Jugular Vein Grafts with FRAME Mesh in a Porcine Carotid Artery Model

Jaroslav Chlupac^{1, 2}, Jan Frank¹, David Sedmera³, Ondrej Fabian^{4, 5}, Zuzana Simunkova⁶, Iveta Mrazova⁶, Tomas Novak¹, Zdenka Vanourková⁶, Oldrich Benada⁷, Zdenek Pulda⁸, Theodor Adla⁸, Martin Kveton^{4, 9}, Alena Lodererova⁴, Ludek Voska⁴, Jan Pirk¹⁰, and Jiri Froncek^{1, 2, 11}

¹ Transplantation Surgery Department, Institute for Clinical and Experimental Medicine (IKEM), Prague, Czech Republic.

² Department of Anatomy, Second Faculty of Medicine, Charles University, Prague, Czech Republic.

³ Institute of Anatomy, First Faculty of Medicine, Charles University, Prague, Czech Republic.

⁴ Clinical and Transplant Pathology Centre, IKEM, Prague, Czech Republic.

⁵ Department of Pathology and Molecular Medicine, Third Faculty of Medicine, Charles University, and Thomayer University Hospital, Prague, Czech Republic.

⁶ Experimental Medicine Centre, IKEM, Prague, Czech Republic.

⁷ Laboratory of Molecular Structure Characterization, Institute of Microbiology of the Czech Academy of Sciences, Prague, Czech Republic.

⁸ Department of Imaging Methods, IKEM, Prague, Czech Republic.

⁹ Third Faculty of Medicine, Charles University, Prague, Czech Republic.

¹⁰ Cardiovascular Surgery Department, IKEM, Prague, Czech Republic.

¹¹ First Surgical Clinic, First Faculty of Medicine, Charles University, Prague, Czech Republic.

Background: Autologous vein grafts are commonly utilized in cardiovascular bypass surgeries. Despite their frequent use, these grafts often fail due to vein graft disease.

Objective: This study was designed to assess the effects of the latest-generation external FRAME (VGS, Tel Aviv, Israel) support on the remodeling of vein grafts using a porcine preclinical model [1].

Methods: Autologous internal jugular vein interposition grafting was conducted in the carotid arteries of pigs over a one-month period. Out of the total grafts, four were reinforced with FRAME mesh, and seven served as unsupported controls. These grafts were evaluated using flowmetry, angiography, macroscopy, and microscopy.

Results: After one month, the patency rate for the FRAME-supported grafts was 100% (4/4), while it was 43% for the unsupported controls (3/7, Log-rank $p = 0.071$). Angiographic examination at explant revealed that FRAME-supported grafts significantly exhibited more areas with no or mild stenosis (9/12 segments) compared to the controls (3/21 segments, $p = 0.0009$). The blood flow in the FRAME-supported grafts was higher (145 ± 51 ml/min) than in the controls (46 ± 85 ml/min, $p = 0.066$). The area and thickness of neo-intimal hyperplasia (NIH) at the proximal anastomoses were comparable between the FRAME and control groups: 5.79 ± 1.38 mm² vs. 6.94 ± 1.10 mm² ($p = 0.558$), and 480 ± 95 $\mu\text{m}^2/\mu\text{m}$ vs. 587 ± 52 $\mu\text{m}^2/\mu\text{m}$ ($p = 0.401$), respectively. However, the NIH area and thickness in the midgraft portions were significantly lower in the FRAME group compared to the control group: 3.73 ± 0.64 mm² vs. 6.27 ± 0.64 mm² ($p = 0.022$) and 258 ± 49 $\mu\text{m}^2/\mu\text{m}$ vs. 518 ± 36 $\mu\text{m}^2/\mu\text{m}$ ($p = 0.0002$), respectively.

Conclusions: The perivascular FRAME mesh enhanced the patency and reduced the incidence of stenosis in vein-to-carotid artery grafts within our porcine model, particularly in the midgraft regions. Midgraft neo-intimal hyperplasia was substantially less pronounced in the FRAME-supported grafts compared to the controls, demonstrating a two-fold difference in thickness.

[1] Chlupac J., Frank J., Sedmera D., et al. *Biomedicines*. 2024; accepted, manuscript ID 3016779.

Funding sources: Supported by the National Institute for Research of Metabolic and Cardiovascular Diseases (Programme EXCELES, ID Project No. LX22NPO5104) - Funded by the European Union – Next Generation EU.

Financial disclosures: none.

Relaxin Receptor Agonist ML290 Inhibits Calcification in Vascular Smooth Muscle Cells

Ana M Valentín Cabrera¹, Valery Butto¹, Kenneth J Wilson Ph.D.², Mark J Henderson Ph.D.², Juan J Marugan Ph.D.², Joshua D Hutcheson Ph.D.¹, Alexander Agoulnik Ph.D.¹

¹*Department of Biomedical Engineering, Florida International University, Miami, FL, USA* ²*Early Translation Branch, National Center for Advancing Translational Sciences, National Institutes of Health, Bethesda, MD, USA*

Vascular calcification contributes to the rupture of atherosclerotic plaques—the leading cause of heart attacks. No therapeutics exist to treat vascular calcification. We hypothesized that relaxin, a vasoprotective and anti-fibrotic small peptide hormone of the insulin/relaxin family, may provide a therapeutic option for vascular calcification. However, recombinant relaxin has short stability *in vivo*, poor bioavailability, and is expensive to synthesize, limiting its clinical utility for chronic conditions such as vascular calcification. ML290 is a biased allosteric agonist of the human relaxin receptor (hRXFP1), which was previously shown to reduce alkaline phosphatase activity in human aortic vascular smooth muscle after 7 days. This study aimed to determine if ML290 may stop the progression of vascular calcification *in vitro* after chronic exposure to pro-calcific media. HCAVSMCs were cultured for 21 days. The first group was cultured with standard media (DMEM, FBS, Pen/Strep). The second group was cultured in pro-calcific media (LAA, BGP, Dexa). The final groups were cultured with pro-calcific media and 1, 5, and 10 μM of ML290. Media was collected every week. The extracellular vesicles were then collected from the media, and a commercially available colorimetric alkaline phosphatase assay was used to determine differences in alkaline phosphatase activity. After 21 days, the cells were fixed and stained with alizarin red to visualize calcification deposition in the wells. The stain was also dissolved and quantified. Data suggest that relaxin agonism reduces alkaline phosphatase activity and calcium deposition, as shown by decreased alizarin red stain, dose-dependently in HCAVSMCs. This study will elucidate the dose-dependent effects of ML290 on human vascular smooth muscle cells.

This research was supported by The Florida Heart Research Foundation.

A PORCINE MODEL OF CONTROLLED PERIPHERAL ARTERY CALCIFICATION

Jason MacTaggart¹, Jennifer Gamache¹, Alexey Kamenskiy²

¹ Department of Surgery, University of Nebraska Medical Center

² Department of Biomechanics, University of Nebraska Omaha

Peripheral artery disease (PAD) in the femoropopliteal artery (FPA) leads to significant morbidity and mortality. FPA calcification contributes to poor clinical outcomes and greatly increases the risk of amputation. Current treatments for calcific lesions are limited and yield suboptimal results. A large animal model that closely mimics calcific PAD and accommodates human-sized devices could enhance the development of safer and more effective therapies. Our objective was to create a swine model of late-stage arterial calcification to test the efficacy and side effects of intravascular lithotripsy (IVL). To induce lesions, swine received injections of CaCl₂ directly into the media and periadventitial spaces of the abdominal aorta, iliac, femoral, and popliteal arteries using a micro-needle catheter. The injection sites were varied to create eccentric and concentric lesions of differing lengths and patterns, with dosage adjusted to prevent aneurysmal expansion. Adjacent non-calcified arterial segments served as intersubject controls. The lesions were allowed to mature over 30 days. Computed Tomography Angiography (CTA) and Intravascular Ultrasound imaging demonstrated ring-like calcification patterns and no pulsatility. Mechanical testing of excised arteries mirrored the mechanical properties of calcified human FPAs, including characteristic circumferential stiffening. Histological examination revealed calcified arteries that closely resembled human calcified FPAs, exhibiting inflammation, thickened and inflamed adventitia with extensive vasa vasorum, degenerated and collagen-rich media with round-shaped smooth muscle cells, severely degraded elastin in the external elastic lamina, and substantial neointimal hyperplasia composed of dense collagen, cellular debris, and glycosaminoglycans. The acute effects of IVL on calcific lesions included periadventitial hematoma, extensive arterial wall damage with complete separation of the tunica media, exposure of the adventitial collagen to flowing blood, and acute thrombosis in smaller arterial segments. This porcine model effectively replicates key aspects of calcific human PAD and provides a rigorously controllable environment to evaluate calcium-modifying treatment impacts and mechanisms.

Funding: This work was supported in part by the Nebraska Research Initiative and NIH awards HL125736, HL147128, and P20GM152301.

Biomimetic Electrospun Tri-layer Tissue Engineered Heart Valve with Low Calcification and Good Regenerative Ability

Jing Liu^{1*}, Zhihong Wang², Deling Kong²

¹Institute of Biomedical Engineering, Chinese Academy of Medical Sciences, Tianjin 300192, China

²Nankai University, Tianjin 300071, China

*liujing@bme.pumc.edu.cn

Background of the clinical problem: Transcatheter aortic valve replacement (TAVR) technique is rapidly advancing in clinic, however, as it expands to low-risk populations and younger patients (age < 65 years), device durability is becoming a weakness.

Objective: Developing tissue-engineered heart valves (TEHVs) as a potential alternative with low calcification and good regenerative ability.

Methods: A biomimetic Tri-layer tissue-engineered heart valve was constructed with poly (L-lactate-co-ε-caprolactone) (PLCL), gelatin (GEL), hyaluronic acid (HA) and silk fibroin (SF), to simulate the fibrosa, spongiosa and ventricular layer of natural heart valves, respectively. To obtain a scaffold with proper strength and regenerative capacity, we optimized the component ratio of each layer. The physical, mechanical and cytocompatibility properties were tested, calcification deposition and regeneration potential were further evaluated by subcutaneous implantation in rat model. Finally, the hydrodynamic function of the new TAVR device was verified.

Results: The strength of the Tri-layer valve could reach up to 10 MPa, significantly higher than that of the PLCL and Mono-layer groups. Most importantly, *in vitro* calcification related gene expression of valvular interstitial cells (VICs) was down-regulated in TEHVs group compared to calcification-inducing group, and calcification levels of TEHVs in *in vivo* assay were below 0.5 μg/mg. Besides, we found HA in the middle layer was very conducive to rapid cell infiltration and good angiogenesis, which ultimately promoted host tissue regeneration at 8 weeks after implantation. **Conclusions:** In this study, we provide a biomimetic Tri-layer electrospun leaflets with appropriate mechanical strength, low calcification and good regenerative capacity, which has great potential as a TEHV leaflet.

Funding: Natural Science Foundation of China (no. 82272158)

Bilayer small diameter vascular graft with dual anti-calcification property by loading with baicalin and cathepsin S inhibitor

Yanjiao Teng¹, Xiaoqin Guo¹, Shufang Wang², Haojun Fan¹, Shike Hou^{1*}, Qi Lv^{1*}, Jie Shi^{1*}

- a. Institute of Disaster and Emergency Medicine, Tianjin University, Tianjin 300072, China.
- b. Key Laboratory of Bioactive Materials for the Ministry of Education, College of Life Sciences, Nankai University, Tianjin 300071, China.

Abstract

Long-term transplantation of small diameter vascular grafts (SDVGs) is associated with calcification risk, a key factor limiting SDVG development in the clinic. Hence, there is an urgent clinical need to develop new SDVGs with anti-calcification properties. Here, we used the decellularized extracellular matrix (dECM) and poly (L-lactide-co- ϵ -caprolactone) (PLCL) as base materials, baicalin was crosslinked, and cathepsin S (Cat S) inhibitor was loaded to prepare PBC-SDVGs by electrospinning. The vascular grafts crosslinked with baicalin reduced the calcium deposition in simulated body fluid experiments and inhibited vascular smooth muscle cells (VSMCs) switch to synthetic phenotype, downregulating the expression level of osteogenic genes. Vascular grafts containing both baicalin and Cat S inhibitor can effectively promote macrophage polarization toward an anti-inflammatory M2 phenotype. Subcutaneous implantation of the rat for 12 weeks demonstrated that the vascular graft reduced the occurrence of calcification. In summary, vascular grafts crosslinked with baicalin and loaded with Cat S inhibitor effectively reduced the osteogenic phenotype and synthetic phenotype transition of VSMCs, inflammation occurrence, and exhibited dual anti-calcification effects. The dual anti-calcification vascular graft provides a promising strategy for the design and construction of anti-calcification SDVGs with significant clinical application potential.

Acknowledgements

This work was supported by the Scientific Research Translational Foundation of the Tianjin Natural Science Foundation (20JCYBJC01240), the National Natural Science Foundation of China (31870966), the Tianjin University Independent Innovation Fund of China (2021XZS-0025), the Scientific Research Translational Foundation of Wenzhou Safety (Emergency) Institute of Tianjin University (TJUWYY2022008, TJUWYY2022020).

Conflict of interest

The authors declare no conflict of interest.

Self-EmPOWERment - Road to a Self-Powered, Intelligent Vascular Graft

H. Middleton¹, Joana O. Gomes,¹ C.S. Rodrigues,² João O. Ventura,² Helga Bergmeister,³ André M. Pereira,² Inês C. Gonçalves,^{1,4} Andreia T. Pereira^{*,1,4}

¹i3S – Institute for Research and Innovation in Health, University of Porto

²IFIMUP – The Institute of Physics for Advanced Materials, Nanotechnology and Photonics, Department of Physics and Astronomy of the Faculty of Sciences, University do Porto

³CBMRS - Center for Biomedical Research, Medical University of Vienna

⁴INEB – National Institute of Biomedical Engineering, University of Porto

*: mail: andreia.pereira@i3s.up.pt

Keywords: biomaterials, biomedical engineering, energy harvesting, triboelectricity, vascular grafts,

Even after 70 years, vascular grafts (VG) have not seen any of the significant improvements desperately needed to improve patient survival rate[1]. The unexpectedness of graft failure raises the need for continuous monitoring of VG performance, as there are rarely symptoms prior to a life-threatening event. However, this is not yet clinically implemented, since the necessary electronics rely on batteries that are bulky, toxic, and have a short lifespan, requiring replacement surgeries. Recently, it was described that fluids flowing through tubular structures can generate electrical energy, by so-called solid-liquid triboelectric nanogenerators (SL-TENG) [2].

Our main aim is to evaluate if blood flowing through commercially VGs is able to generate electric energy to supply the electronics needed for intelligent-VG.

SL-TENG based on commercially available ePTFE VGs (SL-TENG-ePTFE) were assembled by painting a conductive ink on the outer surface. An endovascular simulator mimicked physiological blood flow through SL-TENG-ePTFE. Open circuit voltage and energy storage potential were evaluated using an electrometer. The best VGs were evaluated *in vivo* by establishing an AV-shunt in pig to promote the blood flow through the SL-TENG-ePTFE.

By testing different SL-TENG-ePTFE *in vitro*, it was possible to generate 1.68 V and charge 200 μ F capacitors up to 2.6 V. The best SL-TENG-ePTFE was able to supply the emission of a Bluetooth signals via a commercially available circuit every 23 min. The capacity to generate energy by blood flowing through vascular grafts was also proved *in vivo*, being possible to obtain 0.50 V in open circuit.

Our straightforward SL-TENG systems show the ability to be used as an endless source of power towards a smart VG that can monitor its performance 24/7 which will, in the future, prevent their unexpected failure that culminates in patient death.

Funding Sources. HORIZON-EIC-2022-PATHFINDERCHALLENGES-01projectBlood2Power (GA101115525) funded by the European Union; FCT through 2022.05030.PTDC; 2021.01807.CEECIND.

References.

1. Ratner, B., *Vascular Grafts: Technology Success/Technology Failure*. BME Front, 2023. **4**: p. 0003.
2. Nam, G.H., et al., *A New Pathway for Liquid–Solid Triboelectric Nanogenerator Using Streaming Flow by a Novel Direct Charge Transfer*. Advanced Energy and Sustainability Research, 2020. **1**(1).

Electrospun Vascular Grafts with Artery-Tuned Anisotropy

Elizabeth Zermeno¹ and Kaspars Maleckis¹¹Department of Biomechanics, University of Nebraska at Omaha

Electrospun elastomeric biomaterials hold high promise in developing novel mechanically biomimetic vascular grafts with improved clinical performance. Controlling fiber alignment in such materials can potentially achieve mechanical anisotropy similar to that of human arteries, further advancing graft performance. However, the level of achievable anisotropy and its relation to microstructure parameters has not been sufficiently studied to allow the development of biomimetic grafts with artery-tuned anisotropy. This work aimed to develop a theoretical model for anisotropy control in electrospun vascular grafts and quantify the level of achievable anisotropy values through a parametric study.

Electrospun elastomeric fabrics with varying levels of fiber alignment were produced through the modulation of rotational collector speed. Fiber alignment was quantified by calculating Chebyshev order parameters from fiber orientation histograms. Experimental anisotropy (A_e) was quantified as the ratio of circumferential and axial stiffness values and theoretical anisotropy (A_t) was obtained by calculating Krenchel's fiber efficiency factor from the orientation histogram in both principal directions.

Our results demonstrate that fiber alignment through modulation of rotational collector speed allows mimicry of a wide range of physiological anisotropy values seen in human arteries. A_t values show good agreement with A_e at varying levels of fiber orientation, validating this theoretical framework. Furthermore, we developed a mathematical relationship, showing that A_t can be calculated directly from Chebyshev order parameters. These results provide new insight into structure-property relationships in aligned fibrous biomaterials and provide a novel theoretical framework for optimizing the anisotropy in electrospun elastomeric vascular grafts. Such grafts may provide better mimicry of healthy artery biomechanics, ultimately leading to improved clinical outcomes.

This work was in part supported by the NSF (Graduate Research Fellowship No. 163312) and the NIH (grants 1P20GM152301 and 1R61HL173890).

High-resolution auxetic cardiac patches as a potential tool to support myocardium after infarct

Felix Pointner¹, Christopher Riedmüller¹, Luis Pichelkastner¹, Martin Stoiber², Helga Bergmeister^{1,3}, Bruno K. Podesser^{1,3}, Karl H. Schneider^{1,3,4}, Marjan Enayati^{1,3}

1 Center for Biomedical Research and Translational Surgery, Medical University of Vienna, 1090 Vienna, Austria

2 Center for Medical Physics and Biomedical Engineering, Medical University Vienna, Austria

3 Ludwig Boltzmann Institute for Cardiovascular Research, Vienna 1090, Austria

4 Department of Biomedical Engineering, Tufts University, Medford, MA 02155, USA.

Introduction: Cardiac patches aid heart function after myocardial infarction (MI), adding strength, supplying growth factors and bridging impaired conduction. Auxetic microstructures, stretching perpendicular to the applied force, can add mechanical properties favorable for a demanding, anisotropic implantation site like the myocardium. Herein we aim to fabricate auxetic cardiac patches of synthetic polymers to be functionalized with composite hydrogels for implantation after MI.

Methods: Micropatterned PCL scaffolds were fabricated on a 3D bioplotter by a custom printing protocol for precise control over printhead movement. Mechanical properties were studied in unidirectional, quasistatic tensile tests. Biocompatibility was assessed via XTT, live/dead assays and SEM. Hemocompatibility was investigated via hemolysis and clot formation assays. Inflammatory response was studied via expression of macrophage and cytokine markers. In a pilot *ex-vivo* study in a Langendorff isolated heart system, patch applicability and effects on cardiac function were evaluated on rabbit and rat heart models.

Results: Optimizing the fabrication process for different polymers resulted in uniform scaffolds with defined pore sizes and tunable mechanical properties. Tensile tests confirmed an anisotropic stiffness-ratio. Ultimate strain (7-20%) was close to species-specific myocardium (15-22%). Biocompatibility was proved in cell viability assays and SEM. Patches showed low hemolysis rate. Immunomodulatory effects of the patches were observed. Pro-inflammatory marker expression was significantly reduced after 1 week (CD80, CCR7, IL-1a & TNF-a). For *ex-vivo* testing, auxetic patch parameters were adjusted to the anisotropic ratio of effective stiffness of the model organ. Patches were adhered by fibrin sealant and sutures, without compromising epicardial tissue or patch movement, resulting in modulation of cardiac function.

Conclusion: 3D printing of auxetic, mechanically tunable and biocompatible cardiac patches showed promising results, offering a therapeutic platform for supporting cardiac regeneration, which could be scalable and translatable clinics.

The study was partly funded by the Ludwig Boltzmann Institute for Cardiovascular Research.

Radiation-Modulated Decellularized Pericardium Vascular Patches with Wharton's Jelly Stem Cells in a Porcine Carotid Artery Model

Jan Frank¹, Jaroslav Chlupac^{1,2,3}, Elena Filova³, Ondrej Fabian^{4,5}, Zuzana Simunkova⁶, Iveta Mrazova⁶, Tomas Novak¹, Zdenka Vanourková⁶, Zdenek Pulda⁷, Martin Kveton^{4,8}, Irena Vackova³, Simon Prazak³, Theodor Adla⁷, Alena Lodererova⁴, Ludek Voska⁴, Lucie Bacakova³, and Jiri Froncek^{1,2,10}

¹ Transplantation Surgery Department, Institute for Clinical and Experimental Medicine (IKEM), Prague, Czech Republic.

² Department of Anatomy, Second Faculty of Medicine, Charles University, Prague, Czech Republic.

³ Laboratory of Biomaterials and Tissue Engineering, Institute of Physiology of the Czech Academy of Sciences, Prague, Czech Republic.

⁴ Clinical and Transplant Pathology Centre, IKEM, Prague, Czech Republic.

⁵ Department of Pathology and Molecular Medicine, Third Faculty of Medicine, Charles University, and Thomayer University Hospital, Prague, Czech Republic.

⁶ Experimental Medicine Centre, IKEM, Prague, Czech Republic.

⁷ Department of Imaging Methods, IKEM, Prague, Czech Republic.

⁸ Third Faculty of Medicine, Charles University, Prague, Czech Republic.

⁹ Cardiovascular Surgery Department, IKEM, Prague, Czech Republic.

¹⁰ First Surgical Clinic, First Faculty of Medicine, Charles University, Prague, Czech Republic.

Background: There is a demand for various types of vascular substitutes in cardiovascular and oncological surgery, particularly when autologous grafts are unavailable due to previous use.

Objective: Our aim was to explore preclinically the in vivo performance of tissue-engineered vascular patches.

Methods: The patches were fabricated from the porcine pericardium by decellularization using sodium dodecyl sulfate and DNase. We created four groups: non-irradiated or electron beam (50 kGy) irradiated samples, with or without static recellularization with human Wharton's jelly stem cells (WJCs) attached through fibrin/heparin/VEGF assemblies: 1. Non-irradiated + WJC, 2. Irradiated + WJC, 3. Non-irradiated, and 4. Irradiated. The patches (three per group) were implanted in twelve carotid arteries in six pigs for one month and evaluated through flowmetry, angiography, macroscopy, and microscopy. Data presented as mean \pm SE were compared using one-way analysis of variance (ANOVA).

Results: All patched carotids remained patent and were endothelialized without stenosis signs. No significant differences in blood flow were observed across the groups and time points ($p = 0.8178$). Three clinically silent aneurysms developed, each one in a different group (Non-irradiated + WJC, Irradiated + WJC, and Irradiated) and were excluded from further analysis. Microscopical examinations revealed significantly thicker neo-intimal hyperplasia (NIH) in the Non-irradiated + WJC group ($867 \pm 90 \mu\text{m}^2/\mu\text{m}$) compared to the Non-irradiated group ($615 \pm 26 \mu\text{m}^2/\mu\text{m}$, $p < 0.05$). Though significant, this negative impact was not clinically apparent. NIH thickness in the Irradiated + WJC group and the Irradiated group measured 665 ± 76 and $688 \pm 76 \mu\text{m}^2/\mu\text{m}$, respectively. There were no other differences among the groups.

Conclusions/interpretations: Modifications such as irradiation or seeding with WJCs did not enhance the in vivo performance of our decellularized porcine pericardial vascular patches compared to unmodified samples. Moreover, aneurysmal degeneration occurred in some cases. Possible explanations include the small size of the patches, insufficient to reveal clinically significant benefits, a small sample size, and challenges associated with decellularization, recellularization, and the use of xenogeneic cells.

Funding sources: Supported by the National Institute for Research of Metabolic and Cardiovascular Diseases (Programme EXCELES, ID Project No. LX22NPO5104) - Funded by the European Union – Next Generation EU.

Financial disclosure: none.

Modulating *in vivo* Long Term Remodeling Outcomes of a Polyurethane Based, Antithrombogenic Tissue Engineering Vascular Graft via Compliance Matching

Katarina M. Martinet (1), David R. Maestas (1), Keishi Kohyama (1), Kang Kim (1,2) William R. Wagner (1,3,4,5), Jonathan P. Vande Geest (1,2,5)

- (1) Department of Bioengineering, University of Pittsburgh, Pittsburgh, PA, USA
- (2) Vascular Medicine Institute, University of Pittsburgh, Pittsburgh, PA, USA
- (3) Department of Surgery, University of Pittsburgh, Pittsburgh, PA, USA
- (4) Department of Chemical Engineering, University of Pittsburgh, Pittsburgh, PA, USA
- (5) McGowan Institute of Regenerative Medicine, University of Pittsburgh, Pittsburgh, PA, USA

Background: Tissue engineered vascular grafts (TEVGs) are a promising solution to many of the challenges faced in small diameter vascular grafting procedures. Compliance matched TEVGs have demonstrated increased vascular smooth muscle cell presence and decreased proinflammatory markers but demonstrate concerns with long term graft viability due to rapid gelatin degradation.

Objective/hypothesis: This study aims to understand the long-term effects of compliance matching a hybrid gelatin/polyurethane TEVG *in vivo*.

Brief methodological details: CM TEVGs were fabricated via electrospinning pure PESBUU and PEUU:gelatin solutions. Grafts were crosslinked in genipin and mechanically tested on a custom microbiaxial device. TEVGs were then implanted as abdominal aortic grafts in Sprague Dawley rats and were monitored for patency, compliance, and hemodynamics using ultrasound for 6 months (n = 8).

Results: Both CM TEVGs and Hypo TEVGs were successfully matched to their target compliances *in vitro*. *In vivo*, the compliance of CM TEVGs and Hypo TEVGs were found to be $0.0012 \pm 0.0004 \text{ mmHg}^{-1}$ (within native range) and $0.0002 \pm 0.0005 \text{ mmHg}^{-1}$, respectively. CM TEVGs experienced significant drops in compliance loss over the first two months of remodeling Hypo TEVGs experienced no shifts. CM TEVGs experience a significant drop in PSV at month 6. Preliminary 28 day explant histology shows complete degradation of the highest gelatin content layer in all CM TEVG samples with abluminal cellularization. Additionally 2 female rats experienced graft dilation (diameter increase) and increased collagen deposition abluminal.

Conclusions/interpretations: Compliance can be successfully modulated with gelatin inclusion. Upon implant, CM TEVGs and Hypo TEVGs fall within their compliance target which can be measured *in vivo* via ultrasound. Neither of the gelatin containing grafts alter vascular hemodynamics. The remodeling profile of CM TEVGs is more dynamic than those of Hypo TEVGs as noted by shifts in compliance.

Funding sources: NIH R01 (HL157017) and NIH T32 (T32HL076124).

Harnessing programmed cell death to guide *in situ* remodeling of human tissue-engineered matrices

Pascal D. Breitenstein¹, Marcy Martin¹, Sarah E. Motta¹, Maximilian Y. Emmert^{1,2,3,4} and Simon P. Hoerstrup^{1,2}

¹ Institute for Regenerative Medicine, University of Zurich, Zurich, Switzerland.

² Wyss Translational Center Zurich, University of Zurich and ETH Zurich, Zurich, Switzerland.

³ Department of Cardiovascular Surgery, Charité Universitätsmedizin Berlin, Berlin, Germany.

⁴ Department of Cardiothoracic and Vascular Surgery, German Heart Center Berlin, Berlin, Germany.

Presenting author: Pascal Breitenstein, pascal.breitenstein@irem.uzh.ch

In situ tissue-engineered heart valves (TEHVs) using human cell-derived tissue-engineered matrices (hTEMs) offer a promising solution to the limitations of current heart valve prostheses. hTEM is a hybrid biomaterial consisting of an extracellular matrix (ECM) and a polymer, which provides initial mechanical stability and functionality and allows for tissue engraftment, repopulation, and remodeling once implanted into a host. Decellularization is a crucial step in the production of hTEMs, aimed at removing cellular components while minimally impacting the ECM. Currently available detergent-based decellularization protocols are insufficient to effectively reduce all cellular residues. We hypothesize that a programmed cell death-assisted decellularization approach, combined with hTEM functionalization with the secretome from dying cells can enhance the adaptive remodeling capacity of hTEM.

TEHVs derived from hTEMs were decellularized using a programmed cell death-assisted method and characterized by histology, biochemical assays, and mass spectrometry. Additionally, TEHVs were evaluated in an *in vitro* pulse duplicator under pulmonary conditions. The secretome from dying human dermal fibroblasts (hDFBs) was characterized and used to functionalize hTEMs.

Our results demonstrate that the programmed cell death-assisted decellularization method meets the decellularization standards in hTEMs. TEHVs decellularized by this method exhibited good leaflet functionality in the *in vitro* tester for up to one hour under pulmonary conditions. Furthermore, the secretome contains high levels of cytokines that influence cell proliferation and trigger a tissue repair response in THP-1 cells.

In conclusion, using programmed cell death to decellularize and functionalize hTEMs is a promising strategy to reduce potential immunogenic residues and introducing biochemical cues that could enhance and guide remodeling *in situ*.

This project was supported by the Maxi-foundation.

Tissue-engineered vascular grafts utilizing automated decellularization of allo- and xenogeneic pericardial and vascular tissues modified with collagen and stem cells 3D bioprinting

Roman Matejka¹, Jana Matejkova¹, Denisa Kanokova¹, Monika Supova², Jaroslav Chlupac³

¹ Department of Biomedical Technology, Faculty of Biomedical Engineering, Czech Technical University in Prague, Kladno, Czech Republic

² Department of Composites and Carbon Materials, Institute of Rock Structure and Mechanics, Czech Academy of Sciences, Prague, Czech Republic

³ Transplantation Surgery Department, Institute for Clinical and Experimental Medicine (IKEM), Prague, Czech Republic

Cardiovascular surgery still lacks suitable biomaterials for whole vessel bypass surgery and patch repair. These currently possess many shortcomings – limited patency for small calibers, limited remodeling, and risk of thrombosis. We aim to prepare tissue-engineered vascular grafts and patches based on decellularized and lyophilized matrices (pericardium and vessels) followed by recolonization with adipose-derived stromal or Wharton's jelly stem cells. Acellular cadaveric donors or animal tissues are an abundant source of promising biomaterials. However, these tissues must be decellularized to minimize the immunogenic response and then processed to reduce thrombogenicity and improve the healing and remodeling process. Decellularization was performed using our custom-built semi-automated system. This system allows the cyclic change of decellularization agents (SDS, Na-Deoxycholate, DNase) and rinsing water, maintaining a reproducible process. Concentrations, cycle times, and their repetitions were optimized based on residual DNA in the tissue and structural and biomechanical properties. Special chambers were created depending on the tissue (tubular and planar) to maintain homogeneous contact and complete decellularization. Recellularization with stromal/stem cells attracts the ingrowing host cells through a paracrine mechanism, improving vascular remodeling in vivo and tending to suppress the formation of intimal hyperplasia, facilitating remodeling. To ensure homogeneous recolonization, we used the 3D bioprinting and microextrusion. Planar tissues were recolonized using the 3D bioprinting method using porcine collagen with incorporated cells. For coating the inner lumen of tubular tissues, we used microextrusion utilizing custom microcannulas and bioprinting for the outer lumen. These microcannulas contain evenly spaced microchannels in which collagen with cells leaving the inner lumen is homogeneously coated. The prepared substrates were then cultivated in perfusion bioreactors with pressure stimulation, promoting cell proliferation and ingrowing into decellularized tissues. Prepared planar patches and tubular grafts were then implanted into pigs for one month of observation. Cell recolonized replacements accelerated remodeling and substrate resorption; WJCs also have immunomodulatory properties.

This research was funded by the Ministry of Health of the Czech Republic grant No. NW24-08-00064 and NW24J-02-00061.

Go with the flow: on the way of advancing native blood vessels as energy harvesters

Ana I. Mendes¹, Daniela M. Sousa¹, Herbert Middleton¹, Helga Bergmeister^{3,4}, André M. Pereira⁵, João O. Ventura⁵, Andreia T. Pereira^{1,2,#}, Inês C. Gonçalves^{1,2,#}

¹ i3S - Instituto de Investigação e Inovação em Saúde, Universidade do Porto, Rua Alfredo Allen, 208, Porto 4200-180, Portugal

² INEB - Instituto de Engenharia Biomédica, Universidade do Porto, Rua Alfredo Allen, 208, Porto 4200-180, Portugal

³ Center for Biomedical Research and Translational Surgery, Medical University of Vienna, Austria

⁴ Ludwig Boltzmann Institute of Cardiovascular Research, Vienna, Austria

⁵ IFIMUP – The Institute of Physics for Advanced Materials, Nanotechnology and Photonics, Department of Physics and Astronomy of the Faculty of Sciences, University do Porto

Equal contribution

Implantable electronic devices are currently limited by the use of batteries to power them. Such batteries have a finite life, are large (hindering their application in smaller devices), and may heat or leak their content (damaging the surrounding tissues). This raises a need for continuous, sustainable and biofriendly sources of energy to supply these devices.

This work proposes an innovative approach based on a fluid-solid triboelectric nanogenerator (Flu-TENG) that harvests energy from blood flowing through the native vessels, an unexplored energy source so far. To develop the Flu-TENG, different electrodes, including conductive silver inks and a conductive polymer, were assembled by brush painting on human umbilical cord arteries, and evaluated for (i) *in vitro* stability; (ii) impact on the intrinsic biomechanical properties of the native arteries, and (iii) efficiency in generating electrical outputs, using PBS as fluid-phase and pumped in a similar way as the pulsatile flow of blood circulation.

Electrodes were successfully assembled and remained stable after washing. Indeed, no particles were detected from the leachables of coated native arteries, nor significant differences were observed when comparing the optical density of uncoated and coated native arteries leachables. Furthermore, electrodes were confined to the outer surface of arteries, with no infiltration into the lumen, as observed by optical and scanning electron microscopy. Interestingly, silver inks did not significantly affect the arteries biomechanics, while the conductive polymer resulted in an increase of the compliance and the maximal burst pressure. Additionally, an electrical output was generated with PBS flowing in a pulsatile rate similar to blood circulation (60 bpm). Collectively, our findings highlight the potential of exploring blood flowing through the native vessels to power implantable electronic devices.

Funding: Project funded by European Union's Horizon EIC Pathfinder Challenges 2022 under grant agreement no. 101115525.

Fabrication and Mechanical Characterization of Near-Field Electrospun Bioresorbable Vascular Grafts Mimicking the Arterial Extracellular Matrix

Alexandra E. Snyder¹, Jada K. Sandridge¹, Adeline E. Nordmoe¹, Evan N. Main¹, Gary L. Bowlin^{1*}

¹Department of Biomedical Engineering, The University of Memphis, Memphis, TN, USA

***E-mail: gbowlin@memphis.edu**

Cardiovascular disease is a global health problem, characterized by the thickening of arterial walls restricting blood flow. One treatment option is a surgical procedure utilizing an autologous or synthetic vascular graft to bypass or replace the diseased arterial segment. It was hypothesized that by mimicking the arterial extracellular matrix in bioresorbable vascular grafts via near-field electrospinning (NFES), vascular templates could be produced with mechanical properties that mimic the native vasculature more accurately than the current synthetic vascular grafts while maintaining sufficient pore sizes to allow transmural ingrowth of capillaries and the reendothelialization of the graft lumen via sprouting endothelial cells. Polydioxanone vascular constructs with circumferential fiber alignment angles of 15°/75° and 30°/60° (0° representing circumferential fiber alignment) were fabricated using a custom built NFES system. The vascular construct mechanical properties were compared to the saphenous vein (SV) and internal mammary artery (IMA) through longitudinal and circumferential uniaxial mechanical testing, suture retention, and burst pressure evaluations. The results demonstrated that the 15°/75° templates were closest to mimicking the native vessel target properties as compared to the 30°; however, neither of the vascular template designs achieved or exceeded all the target values. For the ultimate tensile strength, both the constructs met the SV value on the circumferential axis (2.61 MPa) and the IMA value on the longitudinal axis (4.3 MPa). In terms of suture retention, the 15°/75° template was the only construct that was in the IMA and SV targeted range of 138-200 gf. Finally, the burst pressure testing results indicated that neither of the vascular constructs achieved the level of the SV or IMA (1600-3196 mmHg), however, the 15° constructs were within the lower error of the SV. In conclusion, with further design modifications, NFES constructs have demonstrated promise as small-diameter vascular grafts by mimicking native arterial architecture and mechanical properties.

Promoting cell adhesion and regeneration in a 3D-printed cardiac patch via the human placenta chorion extracellular matrix

Christopher Riedmüller¹, Felix Pointner¹, Sabrina Rohringer¹, Lukas Weber¹, Luis Pichelkastner¹, Marvin Dötzlhofer¹, Marta Bonora³, Bruno K. Podesser^{1,2}, Helga Bergmeister^{1,2}, Karl H. Schneider¹, Marjan Enayati^{1,2}

¹ Center for Biomedical Research and Translational Surgery, Medical University of Vienna, Austria

² Ludwig Boltzmann Institute for Cardiovascular Research, Austria

³ Center for Medical Physics and Biomedical Engineering, Medical University Vienna, Austria

Introduction:

Cardiac patches provide mechanical support and a platform for tissue regeneration after myocardial infarction. However, the hydrophobic surface of 3D-printed polymeric patches can hinder cellular attachment and regeneration. Integration of human placenta chorion extracellular matrix (hpcECM) could induce higher cell affinity and cell adhesion ligand for cardiac related cells resulting an excellent regeneration.

Methods:

Polycaprolactone (PCL) was used to print porous patches, which were then coated with various concentrations of hpcECM. The coating was assessed using an antibody against fibronectin and surface nano-mechanics measurements. Hydrophilicity was measured. Human umbilical vein endothelial cells (HUVEC), human foreskin fibroblasts (HFF), H9c2 and endothelial progenitor cells (EPCs) were utilized to assess the cell viability, attachment, activation via XTT and live-dead-assays and electron microscopy. An initial cell attachment assay was performed. Immunological behavior was analyzed using PCR to investigate macrophage pro- and anti-inflammatory gene expression. Hemocompatibility was evaluated via hemolysis and blood clot formation assays. The in vitro assays spanned from initial time points up to one week.

Results:

With an increase of the hpcECM concentration on PCL a significant increase of the initial cell attachment has been shown. An increase of cell viability was observed in HUVEC, HFF and differentiated H9c2 cells using XTT and live-dead-assays. Coated constructs showed a higher attachment and cell activation of EPCs compared to uncoated PCL constructs. PCR results revealed a slight upregulation of pro-inflammatory markers after 72 hours followed by high expression of anti-inflammatory IL-10 cytokine after one week confirming immunomodulatory behavior of hpcECM. Hemocompatibility was observed in all analyzed groups.

Conclusion:

HpcECM showed a high potential of as a matrix supporting attachment and activation of cardiac related cells compared to its present competitors such as fibronectin. This improvement could lead to better functionality of cardiac patches, supporting cardiomyocyte regeneration in vivo.

Abstract for the Biennial Meeting of the ISACB in Vienna 2024

The Tissue-Engineered, Pre-Vascularized Blood Vessel Wall Construct

Lucie Bacakova¹, Julia Tomsu¹, Yu-Chieh Wu¹, Lucie Svobodova¹, Andrea Hejdova², David Lukas², Martin Molitor³, Zuzana Tirpakova⁴, Lenka Luptakova⁴, Eva Petrovova⁴

¹ Laboratory of Biomaterials and Tissue Engineering, Institute of Physiology of the Czech Academy of Sciences, Prague, Czech Republic.

² Faculty of Science, Humanities and Education, Technical University of Liberec, Liberec, Czech Republic

³ Department of Plastic Surgery, First Faculty of Medicine, Charles University and Na Bulovce Hospital, Prague, Czech Republic

⁴ University of Veterinary Medicine and Pharmacy in Kosice, Kosice, Slovak Republic

Background: The clinically used vascular prostheses are based only on materials, whether synthetic polymeric (e.g. ePTFE, PET) or biological (e.g. decellularized matrices), and lack the cellular component arranged in the physiological layers of the vascular wall.

Objective: We therefore attempted to reconstruct the *tunica media* and *intima* simultaneously on a polymeric scaffold modified with a collagen hydrogel.

Methods: Electrospun nanofibrous membranes made of degradable polyesters (PLA, PCL) were seeded with human adipose tissue-derived stem cells (ASCs). After reaching subconfluence, ASCs were overlaid with a collagen hydrogel mixed with human umbilical vein endothelial cells (HUVECs). These cell-laden and control cell-free constructs were implanted onto the chicken chorioallantoic membrane (CAM) in an *ex ovo* model.

Results: The ASCs gradually migrated into the collagen hydrogel and, together with the HUVECs, spontaneously formed pre-capillaries. The ASCs that remained in the space between the pre-capillaries were then able to differentiate towards smooth muscle cells. When the pre-capillaries reached the surface of the collagen hydrogel, they exited onto it, and the HUVECs formed a confluent layer on top. When implanted on CAM, the chicken blood vessels showed a slightly higher affinity for the cell-free constructs.

Conclusions/interpretations: A pre-vascularized tissue construct mimicking the *tunica media* and *tunica intima* of the blood vessel wall has been created. This construct can be used as a vascular patch in planar form and as a vascular graft in tubular form. After implantation *in vivo*, it is expected that the pre-capillaries in the construct will connect to the capillaries of the surrounding tissue, which will also grow into the construct and help to endothelialize its luminal surface.

Funding sources: National Institute for Research of Metabolic and Cardiovascular Diseases project (EXCELES Programme, Project No. LX22NPO5104) - funded by the EU - Next Generation EU, the Czech Acad. Sci. (*Praemium Academiae* grant No. AP2202), and P JAC Project No. CZ.02.01.01/00/22_008/0004562 of the MEYS, CR, co-funded by the European Union.

Financial disclosure: none.

Biomimetic tri-layered small-diameter vascular grafts with decellularized extracellular matrix promoting vascular regeneration and inhibiting thrombosis with the salidroside

Yanjiao Teng,¹ Ju He,² Adam C. Midgley,³ Xiaoqin Guo,¹ Shufang Wang,³ Qi Lv,^{1} Shike Hou^{1*} and Jie Shi^{1*}*

¹ Institute of Disaster and Emergency Medicine, Tianjin University, Tianjin 300072, China.

² Vascular Surgery, Tianjin First Central Hospital, Tianjin 300192, China.

³ Key Laboratory of Bioactive Materials for the Ministry of Education, College of Life Sciences, Nankai University, Tianjin 300071, China.

Abstract

Small-diameter vascular grafts (SDVGs) are urgently required for clinical applications. Constructing vascular grafts mimicking the defining features of native arteries is a promising strategy. To realize biofunctional SDVGs, anti-thrombotic and pro-regenerative activities are necessary. Here, we constructed a tri-layered vascular graft with a native artery decellularized extracellular matrix (dECM) mimicking the component of arteries. Natural blood vessels dECM maintain bioactive factor contents conducive to vascular tissue regeneration. The porcine thoracic aorta was decellularized, frozen, and milled to retrieve dECM powders from the differential arterial layers. The intima and media dECM powders were blended with poly (L-lactide-co-caprolactone) (PLCL) and utilized as the inner and middle layers of electrospun vascular grafts, respectively. Pure PLCL was electrospun and used as a strengthening sheath for the outer layer. Salidroside (Sal), an anti-thrombotic agent derived from *Rhodiola rosea* plants, was loaded into the inner layer of vascular grafts to inhibit thrombus formation. The incorporation of dECM significantly promoted human umbilical vein endothelial cell (HUVEC) extension adhesion, proliferation, migration, and tube-forming *in vitro*. *In vivo*, studies demonstrated that PLCL-dECM-Sal SDVGs improved extracellular matrix formation, lumen endothelialization, and smooth muscle layer regeneration, compared to PLCL SDVGs lacking dECM. Incorporating Sal promoted macrophage polarization to the M2 phenotype and attenuated calcification, thereby promoting tissue regeneration. Adding Sal and dECM inhibited thrombosis and promoted vascular tissue regeneration successfully, providing a promising rationale for their implementation in the construction of SDVGs.

Funding sources

This work was supported by the Scientific Research Translational Foundation of Wenzhou Safety (Emergency) Institute of Tianjin University (TJUWYY2022008, TJUWYY2022020), the Tianjin Natural Science Foundation (20JCYBJC01240), the National Natural Science Foundation of China (31870966), Tianjin University Independent Innovation Fund of China (2021XZS-0025).

Conflict of interest

The authors declare no conflict of interest.

Differential neutrophil deposition on synthetic versus biological scaffolds for in situ cardiovascular tissue engineering

V.C. Vetter^{1,2}, A.I.P.M. Smits^{1,2}, C.V.C. Bouten^{1,2}

¹Department of Biomedical Engineering, Eindhoven University of Technology, Eindhoven, The Netherlands

²Institute for Complex Molecular Systems, Eindhoven University of Technology, Eindhoven, The Netherlands

Background

Neutrophils are the most prevalent cell type in blood, yet their role in tissue regeneration is not well understood. Recent studies by us and other groups reveal neutrophils are the first cells to colonize cardiovascular scaffolds upon implantation into the bloodstream. In addition, neutrophils have garnered significant attention recently, including extensive research investigating their interactions with biomaterials, such as the formation of neutrophil extracellular traps (NETs), on various surfaces. However the response of neutrophils to different scaffold materials and its implication on the subsequent course of the in situ regenerative response remain important open questions.

Objective

To investigate the relevance of neutrophil deposition on various tissue-engineered cardiovascular (TECV)scaffolds and their significance for subsequent tissue regeneration.

Methods

Primary human neutrophils were seeded onto electrospun: polycarbonate polyurethane (PCBU), poly(4-hydroxybutyrate (P4HB), polycaprolactone (PCL) and decellularized porcine valves (dPHVs). Neutrophil characterization, activation and NET quantification was performed after 4 and 24 hours. Donor-matched peripheral blood mononuclear cells were seeded onto the neutrophil-conditioned scaffolds to assess their influence on macrophage polarization via ELISA, PCR and IHC.

Results

The data reveal neutrophil deposition on all examined electrospun synthetic scaffolds. The NET formation seems to occur in a binary manner and shows local heterogeneity. In contrast, on dPHV, no NET formation was detected. Additionally, preliminary data shows neutrophil deposition on electrospun biomaterials enhances the pro- or anti-inflammatory macrophage response. PCR and

ELISA analyses are currently being conducted to provide a deeper understanding of the genetic and protein secretion profiles.

Conclusions

Our findings indicate that neutrophils deposition occurs on all electrospun scaffolds, with notable heterogeneity among replicate groups. Interestingly, this phenomenon was not observed with dPHVs. We are currently analyzing whether this difference is due to the synthetic versus biological nature of the scaffold or its shape. Further analysis is ongoing to investigate the impact on the subsequent macrophage response.

Acknowledgement

We gratefully acknowledge the Gravitation Program “Materials Driven Regeneration”, funded by the Netherlands Organization for Scientific Research (024.003.013)

Characterization of the inflammatory potential of different vascular graft materials

Sabrina Rohringer^{1,2,3}, Anna-Maria Schmitt^{1,2,3}, Sophie Johanna Specht^{1,2,3}, Pia Hager⁴, Marjan Enayati^{1,2,3}, Karl Heinrich Schneider^{1,2,3}, Christian Grasl^{2,5}, Stefan Baudis^{6,3}, Heinrich Schima^{2,5}, Bruno Karl Podesser^{1,2,3}, Helga Bergmeister^{1,2,3}

- 1 Center for Biomedical Research and Translational Surgery, Medical University of Vienna, Währinger Gürtel 18-20, 1090 Vienna, Austria
- 2 Ludwig Boltzmann Institute for Cardiovascular Research, Währinger Gürtel 18-20, 1090 Vienna, Austria
- 3 Austrian Cluster for Tissue Regeneration, Donaueschingenstraße 13, 1200 Vienna, Austria
- 4 HeartBeat.Bio, Dr. Bohr-Gasse 7, 1030 Vienna, Austria
- 5 Center for Medical Physics and Biomedical Engineering, Medical University of Vienna, Währinger Gürtel 18-20, 1090 Vienna, Austria
- 6 Institute of Applied Synthetic Chemistry, Technical University of Vienna, Getreidemarkt 9/163, 1060 Vienna, Austria

Clinically available synthetic vascular grafts are well suited for the replacement of large diameter vessels, but fail in small diameter applications due to an insufficient long-term healing process. In particular, excessive inflammation and immune cell activation during the first days of implantation can lead to early stenosis and reduced biomechanical properties caused by fast scaffold degradation. The aim of the current study was to investigate inflammatory reactions to different graft materials in the early healing phase.

The interaction of immune cells with decellularized human umbilical cord arteries (dUCAs), thermoplastic polyurethane (TPU)/TPU-urea (TPU/TPUU) and expanded polytetrafluoroethylene (ePTFE) was investigated by seeding monocytes and neutrophils onto the materials in vitro. The expression of pro- and anti-inflammatory proteins and genes was determined by immunofluorescence staining and RT-qPCR. Autologous, TPU/TPUU and ePTFE small diameter vascular grafts (SDVGs) were implanted into the abdominal aorta of male Sprague-Dawley rats for 24 h or one week (n=5 per time point). Immunohistochemistry and cytokine arrays of the anastomotic areas were performed.

TPU/TPUU and ePTFE grafts induced higher proinflammatory protein and gene expression of immune cells in vitro. Qualitative analysis of immunohistochemical staining of one-week-old implants showed higher anti-inflammatory CD163 expression in the graft wall of autologous implants. Cytokine arrays showed a significant upregulation of proinflammatory chemokines CCL2, CCL3, CCL5, CCL11, CXCL2, CXCL5 and the peptide hormone resistin in anastomotic regions of ePTFE implants after 24 hours, which partially persisted after one week.

In summary, ePTFE induced a significantly higher proinflammatory response than dUCAs and TPU/TPUU in vitro and in vivo. Seeding of immune cells on TPU/TPUU resulted in higher proinflammatory gene expression in vitro, which was not confirmed in vivo.

The study was partially funded by the Ludwig Boltzmann Institute for Cardiovascular Research.

Abstract for the Biennial Meeting of the ISACB in Vienna 2024

Development of Vascular Patches with Reconstructed *Tunica Media* and *Tunica Intima*

Authors: Anna Bock^{1,2}, Irena Vacková¹, Elena Filová¹, Šimon Pražák¹, Lucie Svobodová¹, Jaroslav Chlupáč^{1,3}, Andrea Hejdová⁴, David Lukáš⁴, Alex Bryan⁵, Joel D. Bumgardner⁵, Marcy Zenobi-Wong², Lucie Bačáková¹

Affiliations:

- ¹ Laboratory of Biomaterials and Tissue Engineering, Institute of Physiology of the Czech Academy of Sciences, Prague, Czech Republic
- ² Swiss Federal Institute of Technology Zurich (ETH Zürich), Zürich, Switzerland
- ³ Transplantation Surgery Department, Institute for Clinical and Experimental Medicine (IKEM), Prague, Czech Republic
- ⁴ Faculty of Science, Humanities and Education, Technical University of Liberec, Liberec, Czech Republic
- ⁵ Department of Biomedical Engineering, The University of Memphis, Memphis, TN, USA

Background: Coronary and peripheral artery bypass grafting (CABG and PABG) is a standard treatment for coronary and peripheral artery disease (CAD and PAD), utilizing autologous or synthetic vascular grafts. While autologous grafts possess regenerative and anti-thrombogenic properties, they are limited by availability and donor site morbidity. Synthetic grafts face challenges in small-diameter applications due to thrombogenicity and intimal hyperplasia.

Objective: This study aims to develop planar vascular wall grafts with reconstructed *tunica media* and *tunica intima*, leveraging decellularized tissues, synthetic polymers, and various cell types.

Methods: Decellularized human saphenous veins and horse umbilical blood vessels, along with electrospun polymers (PCL and chitosan) are used to construct vascular patches. The scaffolds were modified (plasma treatment, fibrin network coating, gelatin coating) and seeded with adipose-derived stem cells (ASCs) or Wharton's jelly-derived mesenchymal stem cells (WJSCs), which were differentiated into smooth muscle cells (SMCs). The constructs were then seeded with human umbilical vein endothelial cells (HUVECs). Mechanical stimulation (laminar shear stress and pulsatile pressure) will simulate the *in vivo* environment and support differentiation. *In vitro* assessments will include histology, immunocytochemistry and marker expression analysis.

Results: Initial results are expected to demonstrate successful differentiation of ASCs and WJSCs into SMCs, with appropriate marker expression. Histological and immunocytochemical analysis is anticipated to show a bi-layered structure mimicking the vascular wall components, namely the *tunica media* and *tunica intima*.

Conclusions: This research aims to advance vascular tissue engineering by developing planar vascular patches with reconstructed vessel wall components, with potential future development into (small-diameter) tubular structures.

Funding Sources: Supported by the National Institute for Research of Metabolic and Cardiovascular Diseases project (EXCELES Programme, Project No. LX22NPO5104) - funded by the EU - Next Generation EU, by the Czech Acad Sci. (Mobility Plus project No. UoM-24-02), and also by P JAC Project No. CZ.02.01.01/00/22_008/0004562 of the MEYS, CR, co-funded by the EU.

Engineering bioadhesive polymer scaffolds to enhance extracellular matrix hydrogel vascular graft mechanics

Zuhao Li, Zhen Liu, Deling Kong, [Adam C. Midgley](#)

Key Laboratory of Bioactive Materials for the Ministry of Education, College of Life Sciences, Nankai University, Tianjin, China

Background: Lack of recellularization, tissue regeneration, and integration with host tissues are predominant instigators of small diameter vascular graft (SDVG) shortcomings and inevitable failure. Thus, there is a high demand for bioactive ‘off-the-shelf’ vascular graft alternatives to currently available synthetic grafts. Decellularised extracellular matrix (dECM) from vascular tissues reconstituted into hydrogels provide ideal bioactive niches to support recellularization and tissue regeneration. However, construction of vascular substitutes with these materials suffers from lack of mechanical performance.

Hypothesis: Engineering supporting scaffolds may enhance dECM mechanics and allow the construction of suitable pro-regenerative vascular graft alternatives.

Methods: Wet spinning was used for the fabrication of uniform and aligned poly(ϵ -caprolactone) (PCL) fibers. Various methods such as heat treatment and functionalisation with bioadhesive polydopamine (PDA) were used to assess the enhancement of the scaffold properties. Mold-poured bovine aorta dECM hydrogel encapsulated the fiber scaffold and cross-linking methods were used to optimise mechanical properties.

Results: The wet-spun fibers yielded a uniform and criss-cross polymer scaffold structure with high elasticity and strain resistance. Heat treatment further bolstered strain resistance and coating with PDA facilitated strong adhesion to the surrounding dECM hydrogel, which filled the pores between the scaffold fibers. Photo cross-linking ensured the resultant graft had appropriate mechanical properties to be employed as vascular graft.

Conclusions: This work provides a proof-of-concept strategy of employing polymer scaffolds to support and reinforce bioactive hydrogels, offering a promising solution for the manufacture of ‘off-the-shelf’ tissue engineered vascular grafts.

Funding sources: This work was supported by the National Natural Science Foundation of China (NSFC) Research Fund for Excellent Young International Scientists, grant number 82250610231.

Additively manufactured multimaterials with the elastic properties of human tissue

Erik Kornfellner¹, Markus Königshofer¹, Gunpreet Oberoi^{1,2}, Lisa Krainz¹, Arno Krause¹, Ewald Unger¹, Francesco Moscato^{1,3,4}

- 1) Medical University of Vienna, Austria
- 2) ACMIT GmbH, Wiener Neustadt, Austria
- 3) Ludwig Boltzmann Institute for Cardiovascular Research
- 4) Austrian Cluster for Tissue Regeneration

To create the best possible conditions for practicing surgery, the training material should closely resemble real tissue. Despite the rapid growth in the variety of 3D printing materials in recent years, finding the right material to replicate a specific structure is still challenging. Particularly, replicating gradients in elasticity, color, or surface characteristics is not possible with pure materials. Material jetting enables the simultaneous 3D printing of multiple materials, leading to composite materials with new properties. Our studies have investigated how to combine multimaterials to achieve specific elastic properties and how to predict these properties using finite element method (FEM) analysis to obtain the right mixture for mimicking human tissue.

Using material jetting (Connex3, Stratasys Ltd., Minnesota, USA), commonly used, commercially available materials were 3D printed as composites and mechanically tested. These materials were simulated in silico, with particular emphasis on the interfaces between different material phases.

As a result, various multimaterials have been created with Young's moduli ranging from 100 kPa to 2.5 GPa. This includes the successful replication of the haptic properties of periosteum, which also has a similar feeling when sutured with a needle. However, the typical strain to failure is less than 50%, which limits the ability to replicate the high stretchability of materials such as skin or large vessels. In terms of predictive models, it has been shown that it is critical to account for the non-ideal interfaces between materials, as well as the deviations from the planned geometry that occur during 3D printing. However, the employed FEM models can consistently simulate previously tested materials and accurately predict the properties of modified multimaterials.

The authors believe that this approach will enable the rapid and cost-effective production of a variety of training objects for various surgical procedures, such as suturing blood vessels, using 3D printing.

ISACB 19th Biennial Meeting

Analysis of life quality parameters and circulating biomarkers in a prospective surgical aortic valve replacement study.

Authors:

Frieda Maria KAINZ^{1, 2}, Lukas RESEMANN², Michael SORITZ¹, Ouafa HAMZA¹, Elisabeth DUNKEL², Casey ZACHARIAS², Dominik WIEDEMANN², Attila KISS¹, Peter POKREISZ¹, Bruno K. PODESSER^{1, 2}

Affiliations:

1 - Center for Biomedical Research and Translational Surgery, Medical University of Vienna, Vienna, Austria

2 - Department of Cardiac Surgery, University Clinic St. Pölten and Karl Landsteiner University of Health Sciences, Sankt Pölten, Austria

Introduction:

Surgical aortic valve replacement (SAVR) is followed by reverse remodeling of the myocardium and comprises complex physiological and structural adaptation of the heart after cessation of pressure overload. Before operation patients often experience symptoms such as reduced activity, dyspnea, angina, and dizziness. We aimed to reveal the impact of SAVR on life quality and circulating biomarkers reflecting cardiac damage, myocardial stretch, inflammatory status, kidney, and liver function up to one year after SAVR.

Methods:

This prospective single center study was performed on patients older than 18 years of both sexes undergoing elective SAVR with either bioprosthetic or mechanical aortic valves alone or in combination with Bentall-de Bono procedure. Inclusion was based on standard criteria of aortic stenosis. Patients with documented prior myocardial infarction, endocarditis, states necessitating cardiac bypass surgery, electrophysiological device implantation or resuscitation, glomerular filtration rate (GFR) below 60 mL/min/1.73 m², or evidenced oncologic malignancies were excluded. Heart failure functional scoring was performed according to the recommendations of the New York Heart Association (NYHA) and angina according to the Canadian Cardiovascular Society (CCS) scoring. Kansas City (KCCF) and SF-12 life quality questionnaires were collected.

Apart of relevant echocardiographic data, we conducted extensive blood sampling with differential cell count and evaluated the biomarkers reflecting the cardiomyocyte damage (high sensitivity troponin T, HsTnT), general muscular damage (creatinine kinase, CK) myocardial stretch (N-terminal B-type natriuretic peptide, NT-proBNP), systemic inflammation (C-reactive protein, CRP), kidney function (creatinine; BUN; GFR) and liver metabolism (alanine aminotransferase – AST; aspartate aminotransferase - ALT, gamma-glutamyl transferase GGT) prior to surgery, 4-6 days, 6 month and 1 year after the SAVR.

Results:

We included 95 patients, 15 were lost to the follow up and the observed cohort of 80 comprised 14 females and 66 males (mean age: 69±6 and 63±8 years). We evaluated 71 patients at 6 months and 37 after 1 year. SAVR significantly reduced the maximal ascending aortic flow both in male and female patients (from 4.3±0.9 to 2.3±0.4 m/s and from 4.6±1.2 to 2.3±0.2 m/s, respectively).

After SAVR, NYHA and CCS scores decreased significantly. In the NYHA-I their ratio increased from 28 % prior SAVR to 86% and to 91% after 6 months and 1 year, respectively. At 1 year neither of the patients belonged to the most severe NYHA-III or IV classes (Fig1). The CCS classification developed analogically. KCCF revealed time-dependent increase of life quality 1 year after SAVR. Concomitantly, the SF-12 questionnaire reported an improvement of mental and physical health in all patients compared to the 6 months.

Serum NT-proBNP levels were initially higher and elevated to the greater extent in females early after operation with return to the baseline in both sexes after 1 year (Fig 2). HsTnT and CK were elevated only during the early post-operative phase. AST, ALT, and GGT were higher in men at all timepoints and normalized during the follow up. GFR was higher in men and in some women reached the critically low level at 6 months. Creatinine and BUN were increasing gradually in time-dependently. While leucocytes were higher in men at all timepoints, CRP increased more in women early after SAVR.

Conclusions:

Our findings demonstrate that SAVR induces restoration of flow pattern from the left ventricle to the aorta, improves NYHA heart failure and CCS angina classification in patients of both sexes. The life quality questionnaires highlighted postoperative increase of physical activities associated with necessary weight loss and improvement of mental health for the whole population. However, our study highlights substantial sex differences in adaptation to the restoration of valve function upon surgical intervention. Women are subjected to SAVR at more advanced age, generally they are more prone to kidney failure and exposed to pressure overload longer, which is also reflected by the GFR and stretch-related NT-proBNP response, respectively. Current data warrant detailed analysis of molecular pattern in fluid biopsies including unbiased, high throughput OMICs techniques followed by detailed bioinformatical evaluation and their correlation to the physiologic measurements, clinical outcomes as well as cardiac biomarkers.

Disclosure:

Nothing to disclose

Development of Autologous Tissue-Engineered Artificial Heart Valve for Transcatheter Pulmonary Valve Implantation

Yasushi Sato, Takeshi Terazawa, Yusuke Inoue, Yoshiaki Takewa

Advanced Medical Engineering Research Center, Asahikawa Medical University

Background: Current mechanical valves and xenogenic bioprosthetic valves have limitations in anti-thrombogenicity and long-term durability, especially for young patients. Our research focuses on developing autologous tissue-engineered artificial heart valves (Biovalves) using an "in body tissue architecture" technique. In this method, tissue-forming molds are embedded subcutaneously, and only the connective tissue that forms around the molds becomes the grafts for implantation. Biovalves show promise for patients with congenital heart disease (CHD) who require multiple open-heart surgeries, offering potential tissue regeneration and long-term valve function. The aim of this study was to develop Biovalves with self-expandable stents, implant them in large animals using a transcatheter technique, and evaluate valve function and tissue structure.

Methods: 3D-printed plastic molds with an internalized shape memory alloy stent were embedded subcutaneously in adult goats. After 2-3 months, the molds were removed with the surrounding tissue, isolating heart valve-like tissue (Biovalves) consisting of autologous connective tissue integrating the stent (stent Biovalves). The stent Biovalves were then implanted into goat pulmonary valves. After 6 months of follow-up, the Biovalves were harvested for histological analysis.

Results: The stent and Biovalve tissue exhibited strong adhesion and remained firmly fixed under crimping conditions. Biovalves were successfully implanted with minimal complications, and no significant stenosis or regurgitation was observed. There was no thrombotic adherence, even without anticoagulation. The implanted Biovalve bound smoothly to the native tissue, showing cellular migration and ECM reconstruction.

Conclusion: The stent Biovalve can be implanted with minimal invasiveness, offering the possibility of long-term functionality through tissue regeneration. This innovation is a promising treatment for patients with CHD and offers an attractive alternative to conventional surgical intervention.

Acknowledgment: This research was supported by the Japan Agency for Medical Research and Development (AMED) (Grant No. JP23he0422026).

Small Biohybrid Valves and Scaffolds for Implant Applications

Selina Sonntag¹, Sergio Acosta¹, José Carlos Rodríguez-Cabello², Achim Zipse³, Stefan Jockenhoevel¹, Alicia Fernández-Colino¹

¹ *Department of Biohybrid & Medical Textiles (BioTex), AME-Institute of Applied Medical Engineering, Helmholtz Institute, RWTH Aachen University, Aachen, Germany.*

² *Bioforge Lab, University of Valladolid, CIBER-BBN, Valladolid, Spain.*

³ *optimed Medizinische Instrumente GmbH, Ettlingen, Germany.*

Background

Small valve replacements are essential for pediatric patients with cardiac diseases and elderly patients with conditions like chronic venous insufficiency. These replacements must meet high standards of biocompatibility, mechanical robustness, and haemodynamic performance. However, current options often fall short to meet such demanding requirements.

Objective

In this study, we address such medical need by developing biohybrid miniature valves. We fabricated small valves made of thermoplastic polyurethane (TPU) and elastin-like recombinamers (ELRs). We evaluated biocompatibility, mechanical properties, and haemodynamic performance in vitro to determine their suitability as new prosthetic valves.

Methods

Tubular TPU scaffolds were manufactured by electrospinning. The material properties were determined with a uniaxial stretcher (UniVert, CellScale), SEM, and a burst pressure testing device. For biocompatibility testing of the material, platelet adhesion and activation, and cell viability using endothelial cells were performed (ISO 10993). The results were evaluated with confocal microscopy, SEM, and with a cell viability kit (CellTiter-Blue, Promega). The TPU scaffold was coated with ELRs by dip-coating and miniature valves were created by suturing the tube into an adapted and modified stent scaffold (optimed). The valve was tested in bioreactors for haemodynamic performance (ISO 5840).

Results

The electrospun TPU showed excellent elastic properties with strain values over 300% and maximum stresses exceeding 4 MPa, surpassing the values of native valves. The ELR-coated TPU scaffold withstood burst pressures over 1000 mmHg. Blood compatibility tests revealed lower platelet activation for ELR-coated TPU compared to clinically used ePTFE, and good cell viability in contact with (ELR-coated) TPU. Haemodynamic testing of the assembled valve showed regurgitation fractions below 15% and complete valve closure.

Conclusion

The biohybrid valve made with electrospun TPU and ELR-coating shows promising biocompatibility, mechanical properties, and suitable haemodynamic performance under in-vitro simulated physiological conditions, making it a strong candidate for small valve replacements.

Funding

BMBF project “Optimize”

Small-diameter Vascular Grafts with Tunable Compliance via Melt Electrowriting

Kilian M. A. Mueller,¹ Christina Ahrens,¹ Linda Grefen,^{2,3} Felix Kur,² Christian Hagl,² Petra Mela¹

¹ Technical University of Munich, Germany; TUM School of Engineering and Design, Department of Mechanical Engineering, Chair of Medical Materials and Implants; Munich Institute of Biomedical Engineering; Munich Institute of Integrated Materials, Energy and Process Engineering. E-mail: kilian.mueller@tum.de

² Department of Cardiac Surgery, LMU University Hospital, Munich, Germany

³ DZHK (German Centre for Cardiovascular Research), partner site Munich Heart Alliance, Munich, Germany

Background

Synthetic small-diameter vascular grafts (SDVGs) show high failure rates *inter alia* due to intimal hyperplasia. A mismatch in compliance between graft and native vessel has been identified as major contributing factor to intimal hyperplasia.

Objective

Following the paradigm of *in situ* tissue engineering, we propose a multiscale polymeric graft architecture to obtain off-the-shelf SDVGs with a tunable compliance covering the full range of native vessel characteristics and the capability to transform into living tissue.

Methods

Tubular scaffolds (\varnothing 4 mm) with controlled patterns of various fiber winding angles were additively manufactured from polycaprolactone microfibers via melt electrowriting (MEW) and subsequently dip coated in gelatine. The scaffold architecture was based on a luminal microporous spiral pattern complemented by a macroporous mechanically-defining outer spiral pattern. All grafts were assessed for compliance, suture retention, burst pressure, and kinking behavior following ISO 7198. Grafts were anastomosed to explanted saphenous veins and porcine coronary arteries.

Results

Controlling the fiber winding angle allowed to tune the compliance from $0.6 \pm 0.4 \%$ (100 mmHg)⁻¹ to $12.7 \pm 2.0 \%$ (100 mmHg)⁻¹ ($n = 3$, $i = 4$), which covers the full physiological range of human veins and arteries. All grafts showed adequate burst pressure (1411 ± 280 mmHg, $n = 5$) and suture retention (0.7 ± 0.3 N, $n = 7$). No kinking was observed for bending radii down to 1.5 mm ($n = 3$). Grafts with venous compliance were successfully anastomosed to human saphenous veins and pulsated with the same amplitude when subjected to pulsatile flow. Arterial compliance was evaluated by coronary artery bypass grafting on porcine hearts.

Conclusion

Our biofabrication strategy leverages microscale additive manufacturing to form physiologically relevant SDVGs featuring compliance tailored to that of the native target vessels. Based on a degradable polymeric scaffold architecture this holds the promise for remodeling into a healthy living vessel.

Thrombus Characterization in Abdominal Aortic Aneurysm and Aortic Dissection Models Using Histology and Scanning Electron Microscopy

Niharika Narra,¹ Luke E. Schepers,¹ Cortland H. Johns,¹ Craig J. Goergen¹

1. Weldon School of Biomedical Engineering, Purdue University, West Lafayette, IN

Aortic dissections (ADs) and abdominal aortic aneurysms (AAAs) are common aortic pathologies. AD is characterized as a tear in the intima, causing separation of the layers within the vessel wall, forming a false lumen. AAA is the localized-dilation and weakening of the aortic wall, resulting in expansion and complex flow patterns. While distinct, both frequently develop thrombus within the lumen (AAA) or within the wall (AD). Due to the differences in hemodynamics, we hypothesize that thrombus structure and composition will vary between these two conditions. To examine this, we utilized murine experimental models, including 1) topical elastase- β -Aminopropionitrile for AAA and 2) angiotensin II-infused apolipoprotein E-deficient mice for AD. Using serial ultrasound measurements, we detected vessel expansion in both models and monitored 1) diameter changes and 2) thrombus deposition. Using histology, we quantified the degree and composition of intramural (IMT) and intraluminal thrombus (ILT) in ADs and AAAs respectively. Color segmentation of Movat's pentachrome slides revealed a significantly higher percent area of fibrin ($p < 0.001$) in ILT, while red blood cells (RBCs) were significantly higher ($p < 0.001$) in IMT. Collagen, elastin, and proteoglycans were present in small amounts in both models. Semi-quantitative examination of histology and scanning electron microscopy images showed layers of fibrin in the ILT structure, while the IMT formed a compact network of fibrin and RBCs. Higher amounts of fibrin appear to correlate with a greater degree of organization in the ILT, which may be caused by the active blood flow present in AAAs. This pattern is mostly absent in the IMT due to the recirculation of blood and low wall shear stress in the false lumen of ADs. These data represent an initial effort to understanding how thrombi formation correlates with vessel growth rate and rupture risk to improve patient outcomes for those suffering from AD and AAA.

Vasoreactivity Testing of Tissue-Engineered Vascular Grafts *In Vivo* in an Ovine Model

Authors: Marissa Guo, MD^{1,2}, Tatsu Watanabe, MD¹, Satoshi Yuhara, MD¹, Christopher Breuer, MD¹

¹Nationwide Children's Hospital, Center for Regenerative Medicine

²The Ohio State University, Department of Surgery

Background: Tissue-engineered vascular grafts (TEVGs) may represent a better alternative to the prosthetic biomaterials that are currently used in cardiovascular reconstruction. Composed of a biodegradable scaffold onto which autologous cells may be seeded, TEVGs provide sites for the formation of neotissue when implanted *in vivo*. The resultant neovessel is theorized to have several advantages over prosthetic grafts, including enhanced compliance, potential for growth, and vasoactive functionality. Using an ovine model, we investigated the vasoreactivity of TEVGs implanted as interposition grafts in the inferior vena cava (IVC).

Methods: Three sheep implanted with TEVGs and one implanted with a polytetrafluoroethylene (PTFE) graft were compared to an age-matched control with native IVC. Animals underwent catheterization procedures during which norepinephrine (NE) was injected into the abdominal IVC at doses of 0.25 mcg/kg and 0.75 mcg/kg. Intravenous ultrasound was used to evaluate changes in the cross-sectional area of the mid-graft or mid-thoracic IVC at maximum constriction in response to drug injection.

Results: The average cross-sectional area of TEVGs decreased by 19.0% and 17.6% from baseline following injections of 0.25 mcg/kg and 0.75 mcg/kg of NE, respectively, compared to 23.5% and 67.3% in the native IVC, while the PTFE graft showed no vasoconstriction and minimal change in diameter. Changes in cardiac output (CO) were inversely related to the degree of graft or IVC constriction.

Conclusion: TEVGs more closely resembled native IVC in their response to localized injections of norepinephrine, unlike PTFE grafts, which had no vasoactive function. Vasoconstriction of the IVC or interposition grafts had a direct and immediate effect on cardiac output. The vasoactive function of TEVGs may allow these grafts an enhanced ability to adapt to the body's dynamic physiologic needs.

Funding Sources: This work is supported by the Department of Defense Expansion Award (W81XWH-18-1-0518).

Financial Disclosures: Research is sponsored by Gunze Ltd.

2024 ISACB ABSTRACT**Title**

Sex-dependent Differences in the Progression of Chronic Kidney Disease Induced Cardiac Dysfunction

Authors

Valentina Dargam¹, Anet Sanchez¹, Aashiya Kolengaden¹, Yency Perez¹, Rebekah Arias¹, Ana Valentin¹, Daniel Chaparro¹, Christopher Tarafa¹, Alexandra Coba^{2,3}, Nathan Yapaolo², Emily A. Todd⁴, Monique Williams⁵, Lina A. Shehadeh^{4,5}, and Joshua D. Hutcheson^{1,7}

Affiliations

¹Department of Biomedical Engineering, Florida International University

²Department of Biological Sciences, Florida International University

³Department of Chemistry, Florida International University

⁴Department of Medical Education, University of Miami Miller School of Medicine

⁵Interdisciplinary Stem Cell Institute, University of Miami Miller School of Medicine

⁶Department of Medicine, Division of Cardiology, University of Miami Miller School of Medicine

⁷Biomolecular Sciences Institute, Florida International University

Abstract

Cardiovascular disease (CVD) in chronic kidney disease (CKD) patients exhibits notable sex-dependent differences. Women with CKD have lower incidences of atherosclerotic events and heart failure but higher rates of CVD events in end-stage CKD compared to men. These variations highlight the need for sex-specific CVD diagnosis and management strategies in CKD patients. Electrocardiograms (ECG) can quickly detect cardiac abnormalities, but common ECG markers are often unreliable in CKD patients. In this study, we assess cardiac structural and functional changes during the progression of CKD-induced CVD in mice using electrocardiography and echocardiography to identify how these changes differ due to disease progression and sex.

Adult C57B/6J mice of both sexes were assigned to either a Control group or an Adenine group, which developed CKD and cardiac dysfunction via a high-adenine diet. Mice were sacrificed at 3, 6, 9, and 12 weeks post-CKD induction, and their cardiac structure and function were analyzed using echocardiography and ECG. Differences in cardiac parameters considering sex and regimen type were statistically assessed.

At week 12, left ventricular thickness in Adenine males was 120.18% greater than in Control males and 135.64% greater than in Adenine females, indicating hypertrophy. They also showed higher ejection fraction and prolonged isovolumetric contraction and relaxation times, indicating systolic and diastolic dysfunction. Adenine males had increased QTc interval duration (63.4 ± 10.3 ms) compared to Control males (49.5 ± 1.7 ms) and Adenine females (47.4 ± 4.6 ms), a marker of abnormal repolarization and cardiovascular event risk. We identified that Speak-J interval also differed by sex, with significant increases in Adenine males at week 6 and Adenine females at week 12.

This study demonstrates the feasibility of this mouse model for investigating sex-dependent differences in CKD-induced cardiac dysfunction, which could aid in the development of tailored diagnostic and management strategies for CVD in CKD patients.

Characterization of the immune response and in vivo remodeling of a new generation of allogeneic Cell-Assembled extracellular Matrix after subcutaneous implantation in sheep.

Julien VITRY¹, Fabien KAWECKI¹, Cecile MONFOULET¹, Marie HOURQUES¹, Yoann TORRES¹, Diane POTART¹, Maude GLUAIS¹, Celia GRENET¹, Nicolas L'HEUREUX¹.

¹*Univ. Bordeaux, Inserm, BioTis, UMR1026, F-33000 Bordeaux, France*

The number of bypasses and vascular access procedures for hemodialysis grafts in Europe is increasing. Due to the limited availability of autologous vessels, surgeons often rely on synthetic grafts, which are prone to infection and chronic inflammation. Moreover, synthetic grafts perform poorly for small-diameter applications (< 6mm). A promising therapeutic is tissue-engineered vascular grafts (TEVGs).

We developed a biological TEVG using a cell-assembled extracellular matrix (CAM) and a textile-based approach. Fibroblast cells synthesize CAM sheets in vitro, which are then processed into yarns and woven into vascular substitutes.

To better simulate clinical conditions, we used a sheep model to produce CAM sheets comparable to human-origin sheets. These ovine CAM yarns were implanted in sheep to study the host immune response in an allogeneic context. We compared the host response to various CAM yarns (devitalized vs. decellularized, laser vs. blade cutting, the addition of an anti-inflammatory molecule, and xenogenic vs. allogeneic CAM yarns) to select the best-performing yarns for weaving our TEVG.

The CAM yarns were implanted subcutaneously in sheep and explanted after 2, 4, 12, and 24 weeks to assess changes in tensile strength, inflammatory response, and remodeling through histological analyses and immunofluorescence.

After 24 weeks, all CAM yarns were retrieved and found to be surrounded by fibrous neo-tissue. The tensile strength of the CAM threads increased over this period, indicating an absence of a degrading inflammatory response.

Differences in immune response and remodeling are currently being analyzed to evaluate the impact of decellularization and laser processes. Human CAM yarns showed a similar increase in strength, suggesting low immunogenicity of our CAM material. Based on these results, CAM yarns are compatible with weaving a TEVG that will maintain its structural integrity and mechanical properties in vivo.

This project is funded by an European Research Council grant.

Non-invasive electrophysiological evaluation of new-onset atrial fibrillation after cardiac surgery: Preliminary results from the BigMap Study

David Santer^{1,2,3}, Simon Adrian Amacher⁴, Brigitta Gahl¹, Jules Miazza¹, Luca Koechlin^{1,5}, Denis Berdajs^{1,2}, Nadine Cueni⁴, Michael Kühne^{2,5,6}, Christian Mueller^{2,5,6}, Stefan Osswald^{2,5,6}, Oliver Reuthebuch^{1,2}, Christian Sticherling^{2,5,6}, Martin Siegemund^{2,4,6}, Friedrich Eckstein^{1,2}, Alexa Hollinger^{2,4}, BigMap Investigators

1 Department of Cardiac Surgery, University Hospital Basel, Basel, Switzerland.

2 Medical Faculty of the University of Basel, Basel, Switzerland.

3 Center for Biomedical Research and Translational Surgery, Medical University of Vienna, Vienna, Austria

4 Intensive Care Medicine, University Hospital Basel, Basel, Switzerland.

5 Cardiovascular Research Institute (CRIB) and Department of Cardiology, University Hospital Basel, University of Basel, Switzerland.

6 Department of Clinical Research, University of Basel, Basel, Switzerland.

Objective

New-onset atrial fibrillation after cardiac surgery (NOAF) is a frequent complication and is associated with postoperative stroke, increased mortality, prolonged hospital length of stay, and higher treatment costs. The structures maintaining NOAF are mainly unknown. Thus, the present study aims to describe electrophysiological patterns of NOAF that might be future therapy targets.

Design and method

Consecutive patients undergoing cardiac surgery have been included and monitored for the first seven postoperative days for the development of NOAF. Patients with NOAF underwent non-invasive electrophysiological mapping using a 252-electrocardiogram vest to identify focal and rotational potential drivers of NOAF (PDs). After mapping, a computed tomography scan of the chest was performed to generate a 3-dimensional model of the atria, thus allowing an identification of NOAF maintaining structures. The primary outcome was the electrophysiological description of potential drivers (PD) of NOAF and NOAF-maintaining structures within the atria.

Results

Of 205 enrolled patients, 62 (30%) developed NOAF. Electrophysiological mapping was performed in 23 NOAF patients (37%). A median of 29 (21-48; IQR) rotational PDs and a median of 26 (20-35[IQR]) focal PDs were identified per patient. The most frequent localizations of rotational drivers were the upper half of the right atrium (23 patients, 100%), the inferior/posterior left atrium (22 patients, 96%), and the left pulmonary veins (22 patients, 96%). The most frequent localizations of focal drivers were the left (22 patients, 96%) and right pulmonary veins (20 patients, 87%) and the upper half of the right atrium (22 patients, 96%).

Conclusions

Based on these preliminary results, structures such as the upper right atrium might play a role in maintaining NOAF and might be targets for future preventive or therapeutic strategies. As the study is ongoing, more precise results with identifying risk factors for specific locations and associations with outcomes are expected.

Financial Disclosures: David Santer has received speaker honoraria, educational grants from Medtronic as well as research grants from Medtronic (Schweiz) AG, Mussler Medical Supply, Freiwillige Akademische Gesellschaft Basel, Mach-Gaensslen-Foundation and Fondation Andreas P. Naef pour la chirurgie thoracique. Simon Amacher has received research grants from Mach-Gaensslen-Foundation and Nora van Meeuwen-Häfliger Stiftung. Jules Miazza has received a research grant from Fondation Andreas P. Naef pour la chirurgie thoracique.

The Solution is Dilution: Parabiosis of WT to LYST Mutant Improves TEVG Performance

Delaney J Villarreal^{1,2}, Mackenzie E Turner^{1,2}, Sahana Rajesh^{1,2}, Jingru Che¹, Joseph T Leland^{1,2}, Tai Yi¹, Christopher K Breuer¹

¹Nationwide Children's Hospital, Abigail Wexner Research Institute

²Ohio State University

Tissue-engineered vascular grafts (TEVGs) for use in the repair of single ventricle disease are limited due to stenosis. We discovered that LYST gene mutation protects against graft stenosis. Global LYST mutation is therapeutically inviable due to numerous cellular dysfunctions. Efforts to design a therapeutic led us to investigate whether 100% of cells need the LYST mutation to have a therapeutic benefit or if partial LYST dysfunction will be beneficial without completely impairing cellular functions. Our hypothesis is that LYST will protect against stenosis at a lower ratio of mutated cells. We used a surgical parabiotic model to determine whether a 50:50 ratio of WT to LYST mutant cells could prevent stenosis. Two mice are surgically attached at the elbow, knee, and skin, allowing formation of shared microvasculature and circulating cells. We investigated 11 parabiotic pairs of WT-GFP and LYST KO mice. Flow cytometry showed that 50% of all live circulating cells were GFP positive and 50% were GFP negative by 6-weeks, indicating equilibration of the pair's cells. Additionally, we injected fluorescent nanoparticles (0.05 um and 2 um) into the WT mouse. All assessed nanoparticles crossed over from one mouse to the other within 10 minutes of injection. We assessed graft patency via explantation at 2-weeks. Of the six assessed pairs, 100% of the WT-GFP mice were patent. This suggests that sharing 50% of the circulating cells was sufficient to protect against TEVG stenosis. Understanding this principle of LYST functionality will be essential when developing therapeutics.

Funding from 1R01HL157491-01, 3R01HL157491-02S1. CKB discloses funding from Gunze Ltd.

Powering the Future: Sputtering Technique for Superior Flu-TENG Performance

Ana Catarina Almeida,^{1,2} Ana L. Pires,³ H. Middleton,^{1,2} Cátia R. S. Rodrigues,³ J. O. Ventura,³ A. M. Pereira,³ Helga Bergmeister,^{4,5} Inês C. Gonçalves,^{1,2} Andreia T. Pereira^{1,2}

¹ i3S - Instituto de Investigação e Inovação em Saúde, Universidade do Porto, Rua Alfredo Allen, 208, Porto 4200-180, Portugal

² INEB – Instituto de Engenharia Biomédica, Universidade do Porto, Rua Alfredo Allen, 208, Porto 4200-180, Portugal

³ IFIMUP - Institute of Physics for Advanced Materials, Nanotechnology and Photonics, Faculty of Science, Porto University, 4169-007 Porto, Portugal

⁴ Center for Biomedical Research and Translational Surgery, Medical University of Vienna, Austria

⁵ Ludwig Boltzmann Institute of Cardiovascular Research

Abstract:

Cardiovascular diseases (CVD) are the leading cause of death, claiming 17.9 million lives each year. To address this critical issue, Cardiovascular Electronic Devices (CEDs) have become essential in both treatment (e.g. pacemakers) and diagnosis (e.g. implantable sensors – Digital Health). However, many CEDs face challenges related to their batteries, which often have short lifespans, large sizes, and the potential for toxic leakage, creating a huge demand for battery-free CEDs. Indeed, increasing the lifespan of batteries by 50% would reduce total costs by 22%, representing savings of 12500€ per patient. Recently, our group discovered that the application of liquid-solid triboelectric nanogenerators (Flu-TENGs) to the biomedical field has great potential to revolutionize the way of harvesting electricity from human body. This method involves the movement of a liquid along a solid surface (e.g., synthetic vascular grafts), which generates opposite charges in both materials and results in charge transfer across the interface. Typically, an electrode is placed on top of the solid-phase to collect the electrons generated on its surface. This work aims to correlate the effect of different electrodes deposited by sputtering with the electric output generated with our Flu-TENG system. Different coatings with different thicknesses (25nm to 300nm) were deposited using a diagonal coating method with a rotation target by sputtering. The results suggest that optimizing the electrode materials and deposition processes can significantly enhance the triboelectric performance of Flu-TENGs, achieving maximal peak-to-peak voltages of 2.93V with well-defined peaks and noise reduction of up to 8%, compared with brush painting. By improving the efficiency of charge transfer, we can achieve higher energy output from Flu-TENG. This advancement will be key towards the development of battery-free CEDs, reducing the frequency of surgical interventions and the associated healthcare costs, powering the future of the biomedical field.

Acknowledgements:

This work was financially supported by Portuguese Funds under FCT Fundação para a Ciência e a Tecnologia through the project 2022.05030.PTDC and 2021.01807.CEECIND.

Durable immunomodulatory nanofiber niche for the functional remodeling of cardiovascular tissue

Tonghe Zhu¹, Yonghang Liu¹, Zhuo Liu², Chen Xu²

¹Multidisciplinary Centre for Advanced Materials, Institute for Frontier Medical Technology, School of Chemistry and Chemical Engineering, Shanghai University of Engineering Science, 333 Longteng Rd., Shanghai 201620, P.R. China. E-mail: zhutonghe89@sues.edu.cn.

²Department of Physiology and Pathophysiology, Shanghai Key Laboratory of Bioactive Small Molecules, School of Basic Medical Sciences, Fudan University, 138 Xueyuan Rd., Shanghai 200032, P.R. China.

Functional remodeling and prolonged anti-inflammatory responses are both vital for repairing damage in the cardiovascular system. Although these aspects have each been studied extensively alone, attempts to fabricate scaffolds that combine these effects have seen limited success. In this study, we synthesized salvianic acid A (SA, danshensu) blocked biodegradable polyurethane (PCHU-D) and enclosed it within electrospun nanofibers to synthesize a durable immunomodulatory nanofiber niche (DINN), which provided sustained SA release during inflammation. Given its excellent processability, mechanical properties, and shape memory function, we developed two variants of the DINN as vascular scaffolds and heart patches. Both these variants exhibited outstanding therapeutic effects in in vivo experiments. The DINN was expertly designed such that it gradually decomposes along with SA release, substantially facilitating cellular infiltration and tissue remodeling. Therefore, the DINN effectively inhibited the migration and chemotaxis of inflammatory cells, while also increasing the expression of angiogenic genes. As a result, it promoted the recovery of myocardial function after myocardial infarction and induced rapid reendothelialization following arterial orthotopic transplantation repair. These excellent characteristics indicate that the DINN holds great potential as a multifunctional agent for repairing cardiovascular tissue.




Fig. 1 The scheme of synthetic durable immunomodulatory nanofiber for the functional remodeling of cardiovascular tissue.

Key Words: inflammation; nanofiber niche; cardiovascular tissue; matrix remodeling

Acknowledgements: National Key Research and Development Program of China (2023YFC2412400, 2023YFC2412403)

References

- [1] Fu J, Ding X, Stowell C E T, *et al.* Slow Degrading Poly(glycerol sebacate) Derivatives Improve Vascular Graft Remodeling in a Rat Carotid Artery Interposition Model[J]. *Biomaterials*, 2020, **257**: 120251.
- [2] Lin X, Liu, Y Bai A, *et al.* A viscoelastic adhesive epicardial patch for treating myocardial infarction [J]. *Nature Biomedical Engineering*, 2019, **3**: 632-643.

Development of a simple same-day implantation technique for allogeneic *in vivo* tissue-engineered vascular grafts to be completed in the operating room

Keiichi Kanda (1), Tatsuro Gondai (1), Masashi Yamanami (1), Hidetake Kawajiri (1), Tomoya Inoue (1), Satoshi Gojo (2), Shinichiro Oda (1).

1. Department of Cardiovascular Surgery, Kyoto Prefectural University of Medicine, Japan;
2. Department of Regenerative Medicine, Kyoto Prefectural University of Medicine, Japan

Background of the clinical problem

We have developed the autologous connective tissue vascular grafts for children constructed in subcutaneous spaces of the patients. In 2014, we clinically applied this technology in pediatric pulmonary artery patch augmentation. However, the formation of reliable connective tissue membranes is an issue in high-risk pediatric patients due to limited subcutaneous areas and insufficient regeneration activities.

Objective/hypothesis

In this study, we explored the option of graft creation in healthy parents for allogeneic transplantation to their children. Furthermore, simplification and shortening of the decellularization process are indispensable to achieve same-day transplantation in the operating room. This study focuses on optimizing decellularization process and conducting preliminary animal implantation experiments.

Brief methodological details

Silicone rod molds were implanted subcutaneously in beagle dogs for four weeks, after which the formed connective tissue tubes were excised. These tissue tubes were decellularized using a 1% sodium lauryl ether sulfate (SLES) solution with horizontal shaking (2h/1h/30min). Following decellularization, DNA quantification and tensile strength measurements in the short-axis direction were performed. The tissues were then trimmed into sheets and transplanted as allogeneic patches into another beagle dog's carotid artery. Post-implantation assessments were conducted using ultrasound, and the grafts were excised after three months.

Results

For one hour or more decellularization was required to ensure the complete removal of cellular components from the connective tissue membranes. Tensile strength measurements indicated no significant differences before and after decellularization. During three month-implantation, the grafts did not develop aneurysmal dilation. Morphological examination exhibited no thrombus formation on the luminal surface, which was covered with smooth neointima.

Conclusions/interpretations

Shaking methods successfully simplified and shortened the decellularization process compared with the previous perfusion methods. The decellularized connective tissue membranes maintained mechanical properties and excellent regenerative performance comparable to previous autografts, which suggests their potential as substitutes for allogeneic vascular grafts.

Funding sources

This study was supported by the Japan Society for the Promotion of Science (JSPS) KAKENHI grants JP19H03742, 20H03767, 20K09151, 21K12649 and 21K16499 from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan.

Financial disclosures

Nothing to disclose.

TISSUE ANALYSIS CORE (TAC) CAPABILITIES AT THE CENTER FOR CARDIOVASCULAR RESEARCH IN BIOMECHANICS (CRIB)

Anastasia Desyatova and Alexey Kamenskiy¹

¹ Department of Biomechanics, University of Nebraska Omaha

Supported by the NIH's Center of Biomedical Research Excellence (COBRE) program, the University of Nebraska Omaha has launched the Center for Cardiovascular Research in Biomechanics (CRiB). This center focuses on translational research to develop innovative materials and devices for treating cardiovascular diseases. A key facility, the Tissue Analysis Core (TAC), leverages Nebraska's biomechanics expertise of comprehensive mechanical, structural, and biological evaluations of arteries from >1,000 human subjects aged 12-99 years. TAC has developed protocols and established partnerships with organ procurement organizations to access scarce human tissues, enhancing the scope of research possibilities. TAC supports CRiB's initiatives and offers its specialized services to the broader scientific community. These services include the mechanical assessment of soft tissues and vascular repair materials through multiaxial tensile testing and pulsatile flow circuits. These tests are critical for assessing tissue compliance, performing constitutive modeling, and studying the impact of various risk factors on tissue elasticity. Equipped with a large micro-CT imaging device, TAC provides high-resolution, non-destructive 3D imaging to examine the structural integrity of biological tissues, engineered materials, and medical devices. Additionally, TAC is equipped for Scanning Electron Microscopy, Duplex and Intravascular Ultrasounds, histological evaluations, and medical device prototyping and design, including implantable devices. Collaborating closely with the neighboring University of Nebraska Medical Center, TAC plays a vital role in the preclinical evaluation of devices and materials using a swine model. This collaboration extends to assisting with regulatory compliance, instrument preparation, animal surgeries, and result evaluations. Current projects at CRiB include developing peripheral stents, aortic stent-grafts, peripheral bypass grafts, and wound dressings, all utilizing large animal models. We invite the scientific community and the biomedical device industry interested in human artery analysis and preclinical evaluations to engage with CRiB and TAC. For more information or to discuss potential projects, please contact us at crib.center@unomaha.edu and crib.tac@unomaha.edu.

Funding: P20GM152301.

Guidelines:

Abstracts should include:

Title
All author names (underline the presenting author)
All author affiliations, denoted by number
Main text

Abstract main text should include:

Background of the clinical problem
Objective/hypothesis
Brief methodological details
Results
Conclusions/interpretations
Funding sources
Financial disclosures relevant to the abstract, if any

Format: 11 pt Times New Roman font, single spaced

Limit of 300 words

No figures are allowed

Demographics and clinical relevance of animal models in haemodialysis research: a systematic review

Joost C. de Vries¹, Kimberley E. Wever², M. Nangaku³, Marianne C. Verhaar¹, Karin G.F. Gerritsen¹, Merle M. Krebber¹

Affiliations

¹Department of Nephrology & Hypertension, University Medical Centre Utrecht, The Netherlands

²Radboud University Medical Center, Radboud Institute for Health Sciences, department of Anesthesiology

³Division of Nephrology and Endocrinology, The University of Tokyo, Bunkyo City, Tokyo, Japan

Background & Aims: With the dialysis population growing, improving haemodialysis (HD) is crucial. While innovations undergo extensive preclinical testing, we lack consensus on which animal models or methods of inducing kidney failure are appropriate and on how we should validate dialysis efficiency. We systematically reviewed the literature for kidney failure animal models investigating the effect of dialysis [HD, haemofiltration (HF), HdiaF and ultrafiltration, excluding peritoneal dialysis] innovations on kidney function.

Methods: The methodology was registered on PROSPERO-CRD42022307144. PubMed and Embase were searched for studies detailing a dialysis intervention in any animal model with acute (AKI) or chronic kidney failure. Primary outcomes included reduced kidney function and dialysis-related urea clearance. Other parameters related to species, kidney injury induction and vascular access. We aimed to identify influential factors by meta-regression analysis.

Results: We included n=51 publications reporting on n=72 experimental comparisons. Dialysis was most frequently performed on dogs (61%), rats (20%) or on goats, pigs, cats or sheep. Characteristics such as sex and age were poorly reported (38% and 19%). Studies primarily surgically induced AKI (91%), validated by plasma urea and creatinine increases. Vascular access in dogs and goats was mostly arterio-venous, while veno-venous in pigs and rats. Most studies investigated HD or HF (>95%), with animals undergoing a single dialysis session (median, [IQR 1,0 – 2,0]). Randomization and blinding was poorly reported (<25%). Moreover, there was limited monitoring of key quality indicators including blood pressure and body temperature (<40%) during dialysis. Data synthesis is ongoing

Conclusion: Our systematic review shows low internal validity of studies due to poor reporting and unclear risk of bias. The infrequent use of sepsis or chronic disease models to test dialysis moreover contributes to low external validity. We recommend a better alignment of experimental set-ups with the patient-population in need of dialysis to propel innovations.

Funding: This research is financially supported by ZonMW, MKMD Synthesis of Evidence, Grant [114024171] to MK.

Advancements and Challenges in Cardiovascular Health and Translational Innovation in Bosnia and Herzegovina

Daria Ler¹, Šekib Sokolović², Snežana Brđanović², Hana Hadžić¹, Rasim Jusufović², Roland Fasol²

¹ ASA Institute for Research, Development and Innovation, Sarajevo, Bosnia and Herzegovina

² ASA Hospital, Sarajevo, Bosnia and Herzegovina

Background: Cardiovascular diseases (CVD) remain a leading cause of mortality in Bosnia and Herzegovina, driven by high prevalence rates of risk factors such as hypertension, obesity, and smoking. This clinical problem is compounded by the evolving healthcare landscape and the need for innovative technological interventions and innovations. **Objective:** This study aims to analyze the latest data on cardiovascular health and translational innovation in Bosnia and Herzegovina, evaluate the impact of innovative health technologies, and identify ongoing research efforts and public health initiatives aimed at improving cardiovascular outcomes. **Methodology:** A comprehensive review of recent studies, healthcare reports, and public health data was conducted. Sources included national health statistics, research publications, and reports from international health organizations. The focus was on the prevalence of CVD, risk factors, and the implementation of telemedicine and other digital health solutions. **Results:** Recent statistics indicate that diseases of the circulatory system account for approximately 52.8% of all deaths in Bosnia and Herzegovina, with ischemic heart diseases and cerebrovascular diseases being predominant. Innovative technologies, such as telemedicine and electronic health records (EHR), are gradually being integrated into the healthcare system, improving access to care and patient management. Research efforts are increasingly focusing on local risk factors and the development of tailored interventions. Public health initiatives are promoting lifestyle changes and early detection of CVD. **Conclusions:** Despite the high burden of cardiovascular diseases, the adoption of innovative health technologies and targeted research initiatives are promising steps towards improved management and prevention. Continued investment in healthcare infrastructure and public health campaigns is crucial for mitigating the impact of CVD in Bosnia and Herzegovina. Collaboration with international institutes and adherence to global health standards will further enhance these efforts.

GO-Graft: From Hydrogel to Anti-Adhesive Vascular Graft for Bypass Surgery

Salomé Luís¹, Duarte Moura¹, Kilian Mueller³, Alexander Löwen², Alicia Fernández-Colino², Jizhen Zhang^{4,5}, Joselito Razal^{4,5}, Petra Mela³, Cristina Barrias¹, Fernão Magalhães⁶, Helga Bergmeister⁷, Inês Gonçalves¹, Andreia Pereira^{*1}

¹i3s – Instituto de Investigação e Inovação em Saúde; INEB – Instituto Nacional de Engenharia Biomédica, ²BioTex – Department of Biohybrid & Medical Textiles, Aachen Univ., ³MMI – Medical Materials and Implants, Technical Univ. of Munich, ⁴Institute for Frontier Materials, Deakin Univ., ⁵Institute for Frontier Materials, Deakin Univ.; Manufacturing BU, The Commonwealth Scientific and Industrial Research Organisation; ⁶LEPABE – Lab. of Process, Environmental, Biotechnology and Energy Engineering, Univ. of Porto, ⁷Center for Biomedical Research and Translational Surgery, Univ. of Vienna

Keywords: Vascular Graft, Graphene Oxide, Suture Retention

Heart and limbs small-arteries blockage can lead to death or limb amputation, respectively. Autologous grafts (patient's vessels) are the gold-standard to restore blood flow, but are limited and, while superior than ePTFE-commercial grafts, present thrombosis risk. Thus, we created the GO-Graft, a synthetic, anti-adhesive, small-diameter vascular graft made of FDA-approved hydrogel poly(2-hydroxyethylmetacrylate)(pHEMA) reinforced with graphene oxide (GO), one of Earth's strongest materials. GO-graft prototype overperforms ePTFE in acute thrombogenicity studies in pigs, but cannot be sutured to host arteries.

This work aims to increase its suture retention (SR) from 55 to ≥ 140 g, making GO-Graft viable for patients.

GO-graft 2D films (pHEMA/GO) were produced by *in-situ* polymerization of HEMA monomers with GO and tetraethylene-glycol-dimethacrylate (molding). To improve SR, we explored: incorporation of co-polymers (alginate); fibers (polypropylene(PP) or GO); films(GO); and meshes(polycaprolactone melt-electrowriting (PCL-MEW) or polyethylene terephthalate(PET) textile) before pHEMA/GO polymerization. GO properties were assessed by XPS and TEM; ELR methacrylation was verified by NMR and FTIR. Hydrogels' surface roughness was evaluated through SEM; SR was assessed by tensile tests (ISO-7198). Among the tested conditions of co-polymer incorporation, only 0.03%(w/v) alginate ionically-crosslinked with 0.5–1M Ca^{2+} / Ba^{2+} / Sr^{2+} incorporation increased SR (up to 157g), but the presence of divalent-ions induced thrombus formation, excluding its suitability for GO-graft. PP fibers improved SR (~170g), yet, poor interaction with pHEMA/GO's matrix led to fiber detachment, resulting in unstable films. To improve hydrogel's matrix interaction, GO fibers or films were incorporated, but the slight increase in SR(~80g) was still substandard. The suture retention strength of pHEMA/GO with polycaprolactone melt-electrowriting (PCL MEW) meshes was 137 g, whereas Polyethylene terephthalate (PET) textiles incorporation resulted in an unrupturable hydrogel under the tested conditions. PET meshed incorporated pHEMA/GO is a suturable hydrogel, paving the way for GO-Graft's future application as a vascular graft.

Funding: La_Caixa_Foundation_CI22-00285,
FCT_(2021.01807.CEECIND_&_2022.05030.PTDC) EU for HORIZON-EIC-2022-
PATHFINDERCHALLENGES-01 #101115525

Graphene Oxide: A Masterpiece for Enhancing Decellularized Tissue Applications in Cardiac Regeneration

Andreia T. Pereira^{1*}, Karl H. Schneider^{2,3}, Joana Santos¹, Rubén Pereira¹, Inês P. Fernandes¹, Sabrina Rohringer^{2,3}, Patrícia C. Henriques¹, Christian Grasl⁴, Sofia F. Melo¹, Herbert Kiss⁵, M. Cristina L. Martins¹, Helga Bergmeister^{2,3}, Inês C. Gonçalves¹

¹INEB/i3S - Instituto de Engenharia Biomédica, Portugal;

²Center for Biomedical Research and Translational Surgery, Medical University of Vienna, Austria;

³Ludwig Boltzmann Institute of Cardiovascular Research, Vienna, Austria;

⁴Center for Medical Physics and Biomedical Engineering, Medical University of Vienna, 1090 Vienna, Austria,

⁵Department of Obstetrics and Gynecology, Division of Obstetrics and Feto-Maternal Medicine, Medical University of Vienna, 1090 Vienna, Austria

The development of small-diameter vascular grafts (diameter <5 mm) and effective cardiac regeneration biomaterials is crucial for cardiovascular diseases, the leading cause of death worldwide. Decellularized tissues and gels have promising features for tissue regeneration but weak mechanical properties, limiting their use in load-bearing applications. We propose incorporating graphene oxide (GO) into decellularized extracellular matrix gels (dECM-gels) and decellularized arteries to overcome these limitations and enable their broader application.

Umbilical cord arteries were decellularized and perfused with a GO suspension (C/O ratio 2:1) with ~1.5 µm lateral size to coat the lumen [1]. The dECM gels were obtained from placental tissue and digested with pepsin to obtain pre-gels, which were mixed with different amounts of GO (1-4% v/v) [2]. Gelation was promoted overnight at 37°C.

For decellularized umbilical cord arteries GO-coatings improved their mechanical properties, increasing the maximum force by 27%, burst pressure by 29%, strain by 25%, and compliance by 10%, comparable to human saphenous veins and mammary arteries, used in bypass surgery. Importantly, GO coatings did not compromise endothelial cell adhesion while reducing platelet and bacteria adhesion, potentially preventing thrombosis and infection until re-endothelialization occurs.

In dECM-gels, SEM images showed that GO intercalated within collagen fibers, and rheological tests revealed a remarkable increase in the complex modulus by 21768% at the highest GO concentration. This enables the production of a stable filament for 3D-printing of scaffolds. Although GO did not decrease bacterial adhesion due to its non-exposure on the surface, it reduced clotting time, indicating increased pro-coagulant properties

Overall, GO significantly enhances the mechanical properties of decellularized arteries and dECM-gels and biocompatibility, representing a major advancement in the application of decellularized tissues for cardiac regeneration and as small-diameter vascular grafts for bypass surgery.

References

1 DOI:10.1021/acscami.1c04028 | 2 DOI:10.1186/s40824-023-00431-5

Funding

FCT(2021.01807.CEECIND and 2022.05030.PTDC) EU for HORIZON-EIC-2022-PATHFINDERCHALLENGES-01 GA101115525